

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/98327 A2

- (51) International Patent Classification: **C07K 9/00** (74) Agent: **VIKSINNS, Ann, S.**; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).
- (21) International Application Number: **PCT/US01/13996**
- (22) International Filing Date: **1 May 2001 (01.05.2001)** (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data: **60/213,417** **22 June 2000 (22.06.2000)** **US**
- (71) Applicant (for all designated States except US): **ADVANCED MEDICINE, INC.** [US/US]; 901 Gateway Boulevard, South San Francisco, CA 94080 (US).
- (71) Applicants and
- (72) Inventors: **LINSELL, Martin, S.** [US/US]; 146 West Third Avenue, Apartment 16, San Mateo, CA 94402 (US). **FATHEREE, Paul, R.** [US/US]; 921 Minnesota Street, San Francisco, CA 94107 (US). **JUDICE, J., Kevin** [US/US]; 212 Dolores Street, El Granada, CA 94018 (US). **LEADBETTER, Michael, R.** [US/US]; 335 Beverly Avenue, San Leandro, CA 94577 (US). **ZHU, Yan** [CN/US]; 1021 Lido Lane, Foster City, CA 94404 (US).
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/98327 A2

(54) Title: **GLYCOPEPTIDE CARBOXY-SACCHARIDE DERIVATIVES**

(57) Abstract: Disclosed are glycopeptide derivatives substituted at the C-terminus and/or the R terminus with a substituent that comprises one or more saccharide groups and a carboxy group; and pharmaceutical compositions containing such glycopeptide derivatives. The disclosed glycopeptide derivatives are useful as antibacterial agents.

GLYCOPEPTIDE CARBOXY-SACCHARIDE DERIVATIVES

PRIORITY OF INVENTION

This application claims priority to United States Provisional Application
5 Number 60/213,417, filed 22 June 2000, which application is incorporated
herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

10 This invention is directed to novel saccharide derivatives of glycopeptide
antibiotics and related compounds. This invention is also directed to
pharmaceutical compositions containing such saccharide glycopeptide
derivatives, methods of using such saccharide glycopeptide derivatives as
antibacterial agents, and processes and intermediates useful for preparing such
15 saccharide glycopeptide derivatives.

Background

Glycopeptides (e.g. dalbaheptides) are a well-known class of antibiotics
produced by various microorganisms (*see Glycopeptide Antibiotics*, edited by R.
20 Nagarajan, Marcel Dekker, Inc. New York (1994)). These complex multi-ring
peptide compounds are very effective antibacterial agents against a majority of
Gram-positive bacteria. Although potent antibacterial agents, the glycopeptides
antibiotics are not used in the treatment of bacterial diseases as often as other
classes of antibiotics, such as the semi-synthetic penicillins, cephalosporins and
25 lincomycins, due to concerns regarding toxicity.

In recent years, however, bacterial resistance to many of the commonly-
used antibiotics has developed (*see J. E. Geraci et al., Mayo Clin. Proc.* 1983,
58, 88-91; and M. Foldes, *J. Antimicrob. Chemother.* 1983, 11, 21-26). Since
glycopeptide antibiotics are often effective against these resistant strains of
30 bacteria, glycopeptides such as vancomycin have become the drugs of last resort
for treating infections caused by these organisms. Recently, however, resistance
to vancomycin has appeared in various microorganisms, such as vancomycin-

resistant enterococci (VRE), leading to increasing concerns about the ability to effectively treat bacterial infections in the future (see Hospital Infection Control Practices Advisory Committee, *Infection Control Hospital Epidemiology*, 1995, 17, 364-369; A. P. Johnson et al., *Clinical Microbiology Rev.*, 1990, 3, 280-291; 5 G. M. Eliopoulos, *European J. Clinical Microbiol.; Infection Disease*, 1993, 12, 409-412; and P. Courvalin, *Antimicrob. Agents Chemother.*, 1990, 34, 2291-2296).

A number of derivatives of vancomycin and other glycopeptides are known in the art. For example, see U.S. Patent Nos. 4,639,433; 4,643,987; 10 4,497,802; 4,698,327; 5,591,714; 5,840,684; and 5,843,889. Other derivatives are disclosed in EP 0 802 199; EP 0 801 075; EP 0 667 353; WO 97/28812; WO 97/38702; WO 98/52589; WO 98/52592; and in *J. Amer. Chem. Soc.*, 1996, 118, 13107-13108; *J. Amer. Chem. Soc.*, 1997, 119, 12041-12047; and *J. Amer. Chem. Soc.*, 1994, 116, 4573-4590.

15 Despite the above referenced disclosures, a need currently exists for novel glycopeptide derivatives having effective antibacterial activity and an improved mammalian safety profile. In particular, a need exists for glycopeptide derivatives which are effective against a wide spectrum of pathogenic microorganisms, including vancomycin-resistant microorganisms, and which 20 have reduced tissue accumulation and/or nephrotoxicity.

SUMMARY OF THE INVENTION

The present invention provides novel saccharide glycopeptide derivatives having highly effective antibacterial activity and an improved mammalian safety 25 profile. More specifically, the saccharide glycopeptide derivatives of the invention unexpectedly exhibit reduced tissue accumulation and/or nephrotoxicity when administered to a mammal.

Accordingly, the invention provides a compound of the invention, which is a glycopeptide substituted at the C-terminus and/or the R-terminus with a 30 substituent that comprises one or more (e.g. 1, 2, 3, 4, or 5) saccharide groups and a carboxy (COOH) group; or a pharmaceutically acceptable salt, or stereoisomer, or prodrug thereof. One preferred group of compounds of the invention are glycopeptides substituted at either the C-terminus or the R-

terminus with a substituent that comprises a saccharide and a carboxy group.

Another preferred group of compounds of the invention are glycopeptides substituted at the C-terminus and the R-terminus with a substituent that comprises a saccharide and a carboxy group. Such saccharide/carboxy

- 5 containing substituents can be derived from a dicarboxylic acid or a derivative thereof by coupling one of the carboxy groups with a saccharide.


Preferably, when the glycopeptide is substituted at the C-terminus, the substituent is attached through an amide bond, an ester bond, or a thioester bond.

- Preferably, a nitrogen-linked substituent is attached to the carbonyl group at the
10 C-terminus to form an amide bond. Preferably, the C-terminus substituent comprises one saccharide group and one carboxy group.

Preferably, the glycopeptide is substituted at the C-terminus with a substituent of the formula $-N(R^w)-R^y-R^z$; wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^z is a

- 15 saccharide.

A preferred C-terminus substituent that comprises one saccharide group and one carboxy group is a substituent of formula III:




wherein one of R^g and R^h is a saccharide, and the other of R^g and R^h is OH.

- 20 Preferably, when R^g or R^h is a saccharide it is a N-linked or O-linked saccharide.

Preferably, when the glycopeptide is substituted at the R-terminus, the substituent is attached to the nitrogen of an aminomethyl group that is attached to the R-terminus (i.e. the resorcinol ring). Preferably, the R-terminus substituent comprises one saccharide group and one carboxy group.


- 25 Preferably, the glycopeptide is substituted at the R-terminus with a substituent of the formula $-CH_2-N(R^w)-R^y-R^z$; wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^z is a saccharide.

A preferred R-terminus substituent that comprises one saccharide group and one carboxy group is a substituent of formula IV:




wherein one of R^m and R^n is a saccharide, and the other is OH . Preferably, when R^m or R^n is a saccharide it is a N-linked or O-linked saccharide.

5 Another preferred R-terminus substituent is a substituent of formula V:



Certain glycopeptide derivatives are described in U.S. Patent Application Serial Number 09/470,209, filed 22 December 1999. Accordingly, the compounds of the invention may preferably exclude glycopeptides of formula II:




(II)

wherein: R^{19} is hydrogen and

- a) wherein R^3 is N-(2-amino-2-deoxygluconic acid); R^5 is hydrogen; R^{19} is hydrogen; and R^{20} is $-\text{NH}-\text{CH}_2\text{CH}_2-\text{NH}-(\text{CH}_2)_9\text{CH}_3$; or
- b) wherein R^3 is OH; R^5 is $-\text{CH}_2-\text{N}-(2\text{-amino-2-deoxygluconic acid})$; R^{19} is hydrogen; and R^{20} is $-\text{CH}_2\text{CH}_2-\text{NH}-(\text{CH}_2)_9\text{CH}_3$.

5

A preferred compound of the invention is a glycopeptide of formula I:



(I)

wherein:

- R^1 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic and $-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$; or R^1 is a saccharide group optionally substituted with $-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$, R^f , $-\text{C}(\text{O})\text{R}^f$, or $-\text{C}(\text{O})-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$;

- R^2 is hydrogen or a saccharide group optionally substituted with $-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$, R^f , $-\text{C}(\text{O})\text{R}^f$, or $-\text{C}(\text{O})-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$;

R^3 is $-\text{OR}^c$, $-\text{NR}^c\text{R}^c$, $-\text{O}-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$, $-\text{NR}^c-\text{R}^a-\text{Y}-\text{R}^b-(\text{Z})_x$, $-\text{NR}^c\text{R}^c$,

or

-O-R^c; or R³ is a substituent that comprises a saccharide group and a carboxy group;

- R⁴ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, -R^a-Y-R^b-(Z)_x,
 5 -C(O)R^d and a saccharide group optionally substituted with -R^a-Y-R^b-(Z)_x, R^f,
 -C(O)R^f, or -C(O)-R^a-Y-R^b-(Z)_x;

- R⁵ is selected from the group consisting of hydrogen, halo,
 -CH(R^g)-NR^cR^c, -CH(R^g)-NR^cR^c, -CH(R^g)-NR^c-R^a-Y-R^b-(Z)_x, -CH(R^g)-R^x,
 and -CH(R^g)-NR^c-R^a-C(=O)-R^a, or R⁵ is substituent that comprises a
 10 saccharide group and a carboxy group;

- R⁶ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, -R^a-Y-R^b-(Z)_x,
 -C(O)R^d and a saccharide group optionally substituted with
 -NR^c-R^a-Y-R^b-(Z)_x, or R⁵ and R⁶ can be joined, together with the atoms to
 15 which they are attached, form a heterocyclic ring optionally substituted with
 -NR^c-R^a-Y-R^b-(Z)_x;

R⁷ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, -R^a-Y-R^b-(Z)_x,
 and -C(O)R^d;

- 20 R⁸ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

- R⁹ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

- R¹⁰ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic; or R⁸ and R¹⁰ are joined to form -Ar¹-O-Ar²-, where Ar¹ and Ar² are independently arylene or heteroarylene;

R^{11} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic, or R^{10} and R^{11} are joined, together with the carbon and nitrogen atoms to which they are attached, to form a heterocyclic ring;

R^{12} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic, $-C(O)R^d$, $-C(NH)R^d$, $-C(O)NR^cR^e$, $-C(O)OR^d$, $-C(NH)NR^cR^e$ and $-R^a-Y-R^b-(Z)_x$, or R^{11} and R^{12} are joined, together with the nitrogen atom to which they are attached, to form a heterocyclic ring;

R^{13} is selected from the group consisting of hydrogen or $-OR^{14}$;

R^{14} is selected from hydrogen, $-C(O)R^d$ and a saccharide group;

each R^a is independently selected from the group consisting of alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene;

each R^b is independently selected from the group consisting of a covalent bond, alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene, provided R^b is not a covalent bond when Z is hydrogen;

each R^c is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic and $-C(O)R^d$;

each R^d is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

R^e is a saccharide group;

each R^f is independently alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, or heterocyclic;

R^x is N-linked amino saccharide or an N-linked heterocycle;

X^1 , X^2 and X^3 are independently selected from hydrogen or chloro;

each Y is independently selected from the group consisting of oxygen, sulfur, $-S-S-$, $-NR^c-$, $-S(O)-$, $-SO_2-$, $-NR^cC(O)-$, $-OSO_2-$, $-OC(O)-$, $-NR^cSO_2-$, $-C(O)NR^c-$, $-C(O)O-$, $-SO_2NR^c-$, $-SO_2O-$, $-P(O)(OR^c)O-$,
 5 $-P(O)(OR^c)NR^c-$, $-OP(O)(OR^c)O-$, $-OP(O)(OR^c)NR^c-$, $-OC(O)O-$, $-NR^cC(O)O-$, $-NR^cC(O)NR^c-$, $-OC(O)NR^c-$, $-C(=O)-$, and $-NR^cSO_2NR^c-$;

each Z is independently selected from hydrogen, aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic;

n is 0, 1 or 2; and

10 x is 1 or 2;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof; provided at least one of R^3 and R^5 is a substituent that comprises a


saccharide group and a carboxy group.

Preferably, R^1 is a saccharide group optionally substituted with

- 15 $-R^3-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^3-Y-R^b-(Z)$. More preferably R^1 is an amino saccharide group substituted on the saccharide nitrogen with
- $-CH_2CH_2-NH-(CH_2)_9CH_3$; $-CH_2CH_2CH_2-NH-(CH_2)_8CH_3$;
 - $-CH_2CH_2CH_2CH_2-NH-(CH_2)_7CH_3$; $-CH_2CH_2-NHSO_2-(CH_2)_9CH_3$;
 - $-CH_2CH_2-NHSO_2-(CH_2)_{11}CH_3$; $-CH_2CH_2-S-(CH_2)_8CH_3$;
 - 20 $-CH_2CH_2-S-(CH_2)_9CH_3$; $-CH_2CH_2-S-(CH_2)_{10}CH_3$;
 - $-CH_2CH_2CH_2-S-(CH_2)_8CH_3$; $-CH_2CH_2CH_2-S-(CH_2)_9CH_3$;
 - $-CH_2CH_2CH_2-S-(CH_2)_3-CH=CH-(CH_2)_4CH_3$ (*trans*); $-CH_2CH_2CH_2CH_2-S-(CH_2)_7CH_3$;
 - $-CH_2CH_2-S(O)-(CH_2)_9CH_3$; $-CH_2CH_2-S-(CH_2)_6Ph$;
 - $-CH_2CH_2-S-(CH_2)_8Ph$; $-CH_2CH_2CH_2-S-(CH_2)_8Ph$; $-CH_2CH_2-NH-CH_2-4-(4-$
 - 25 $Cl-Ph)-Ph$; $-CH_2CH_2-NH-CH_2-4-[4-(CH_3)_2CHCH_2]-Ph$; $-CH_2CH_2-NH-CH_2-4-(4-CF_3-Ph)-Ph$;
 - $-CH_2CH_2-S-CH_2-4-(4-Cl-Ph)-Ph$; $-CH_2CH_2-S(O)-CH_2-4-(4-Cl-Ph)-Ph$;
 - $-CH_2CH_2CH_2-S-CH_2-4-(4-Cl-Ph)-Ph$;
 - $-CH_2CH_2CH_2-S(O)-CH_2-4-(4-Cl-Ph)-Ph$; $-CH_2CH_2CH_2-S-CH_2-4-[3,4-di-Cl-PhCH_2O]-Ph$;
 - $-CH_2CH_2-NHSO_2-CH_2-4-[4-(4-Ph)-Ph]-Ph$;
 - 30 $-CH_2CH_2CH_2-NHSO_2-CH_2-4-(4-Cl-Ph)-Ph$; $-CH_2CH_2CH_2-NHSO_2-CH_2-4-(Ph-C\equiv C)-Ph$;
 - $-CH_2CH_2CH_2-NHSO_2-4-(4-Cl-Ph)-Ph$; or
 - $-CH_2CH_2CH_2-NHSO_2-4-(naphth-2-yl)-Ph$. Preferably R^1 is also an amino

saccharide group substituted on the saccharide nitrogen with a 4-(4-chlorophenyl)benzyl group or with a 4-(4-chlorobenzyloxy)benzyl group.

For example, R¹ is preferably a saccharide group of the formula:



wherein R¹⁵ is -R^a-Y-R^b-(Z)_x, R^f, -C(O)R^f, or -C(O)-R^a-Y-R^b-(Z)_x;

5 and R¹⁶ is hydrogen or methyl.

Preferably, R² is hydrogen.

Preferably, R³ is -OR^c, -NR^cR^c, or a nitrogen-linked, oxygen-linked, or sulfur-linked substituent that comprises a saccharide group and a carboxy group.

Preferably, R³ is a substituent of the formula -N(R^w)-R^y-R^x; wherein R^w is
10 hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.

When R³ is not a substituent that comprises a saccharide group and a carboxy group, R³ is preferably -OH; -NH-(CH₂)₃-N(CH₃)₂; *N*-(D-glucosamine); -NHCH(CO₂CH₃)CH₂CO₂CH₃; -NH(CH₂)₃-(morpholin-4-yl); -NH(CH₂)₃-
15 NH(CH₂)₂CH₃; -NH(CH₂-piperidin-1-yl); -NH(CH₂)₄NHC(N)NH₂; -NH(CH₂)₂-N⁺(CH₃)₃; -NHCH(COOH)(CH₂)₃NHC(N)NH₂; -NH-[(CH₂)₃-NH-]₃-H; -N[(CH₂)₃N(CH₃)₂]₂; -NH(CH₂)₃-imidazol-1-yl; -NHCH₂-4-pyridyl; -NH(CH₂)₃CH₃; -NH(CH₂)₂OH; -NH(CH₂)₅OH; -NH(CH₂)₂OCH₃; -NHCH₂-tetrahydrofuran-2-yl; -N[(CH₂)₂OH]₂;
20 -NH(CH₂)₂N[(CH₂)₂OH]₂; -NHCH₂COOH; -NHCH(COOH)CH₂OH; -NH(CH₂)₂COOH; *N*-(glucamine); -NH(CH₂)₂COOH; -NH(CH₂)₃SO₃H; -NHCH(COOH)(CH₂)₂NH₂; -NHCH(COOH)(CH₂)₃NH₂; -NHCH(COOH)CH₂CO₂(CH₂)₃-N⁺(CH₃)₃; -NHCH(COOH)CH₂CO₂(CH₂)₂C(O)-N(CH₃)₂; -NHCH(COOH)CH₂CO₂(CH₂)₃-morpholin-4-yl;
25 -NHCH(COOH)CH₂CO₂(CH₂)₂OC(O)C(CH₃)₃; -NHCH(CH₂COOH)CO₂(CH₂)₃-

- $N^+(CH_3)_3$; $-NHCH(CH_2COOH)CO_2(CH_2)_2C(O)N(CH_3)_2$;
 $-NHCH(CH_2COOH)CO_2(CH_2)_3$ -morpholin-4-yl;
 $-NHCH(CH_2COOH)CO_2(CH_2)_2OC(O)C(CH_3)_3$; $-NHCH(COOH)CH_2CO_2CH_3$;
 $-NHCH(CH_2COOH)CO_2(CH_2)_2N(CH_3)_2$;
 5 $-NHCH(COOH)CH_2CO_2CH_2C(O)N(CH_3)_2$;
 $-NHCH(CH_2COOH)CO_2CH_2C(O)N(CH_3)_2$; $-NHCH(CH_2COOH)CO_2CH_3$;
 $-NH(CH_2)_3N(CH_3)_2$; $-NHCH_2CH_2CO_2CH_3$;
 $-NHCH[CH_2CO_2CH_2C(O)N(CH_3)_2]CO_2CH_2-C(O)-N(CH_3)_2$; $-NHCH_2CO_2CH_3$;
 $-N$ -(methyl 3-amino-3-deoxyaminopyranoside); $-N$ -(methyl 3-amino-2,3,6-
 10 trideoxyhexopyranoside); $-N$ -(2-amino-2-deoxy-6-
 (dihydrogenphosphate)glucopyranose); $-N$ -(2-amino-2-deoxygluconic acid);
 $-NH(CH_2)_4COOH$; $-N$ -(N - CH_3 -D-glucamine); $-NH(CH_2)_6COOH$; $-O$ (D-glucose);
 $-NH(CH_2)_3OC(O)CH(NH_2)CH_3$; $-NH(CH_2)_4CH(C(O)-2-HOOC-pyrrolidin-1$ -
 yl) $NHCH(COOH)-CH_2CH_2Ph$ (*S,S* isomer); $-NH-CH_2CH_2-NH-(CH_2)_9CH_3$; or
 15 $-NH(CH_2)C(O)CH_2C(O)N(CH_3)_2$;

Preferably, R^4 , R^6 and R^7 are each independently selected from hydrogen or $-C(O)R^d$. More preferably, R^4 , R^6 and R^7 are each hydrogen.

Preferably R^5 is hydrogen, $-CH_2-NHR^c$, $-CH_2-NR^cR^e$, $-CH_2-NH-R^a-Y$, $R^b-(Z)_x$, or a substituent that comprises a saccharide group and a carboxy group.

- 20 A more preferred group of compounds of the invention are compounds wherein
 R^5 is $-CH_2-R^p$ wherein R^p is a nitrogen-linked substituent that comprises a
 saccharide group and a carboxy group.

- Preferably, R^5 is a substituent of the formula $-CH_2-N(R^w)-R^y-R^x$; wherein
 R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a
 25 carboxy group; and R^x is a saccharide.

- When R^5 is not a substituent that comprises a saccharide group and a
 carboxy group, R^5 is preferably hydrogen; $-CH_2-N(N-CH_3-D-glucamine)$; $-CH_2$ -
 $NH-CH_2CH_2-NH-(CH_2)_9CH_3$; $-CH_2-NH-CH_2CH_2-NHC(O)-(CH_2)_3COOH$; $-CH_2$ -
 $NH-(CH_2)_9CH_3$; $-CH_2-NH-CH_2CH_2-COOH$; $-CH_2-NH-(CH_2)_3COOH$; $-CH_2$ -
 30 (morpholin-4-yl); $-CH_2-NH-CH_2CH_2-O-CH_2CH_2OH$; $-CH_2-NH-CH_2CH(OH)-$
 CH_2OH ; $-CH_2-N[CH_2CH_2OH]_2$; $-CH_2-NH-(CH_2)_3-N(CH_3)_2$; $-CH_2-N[(CH_2)_3-$
 $N(CH_3)_2]$; $-CH_2-NH-(CH_2)_3$ -(imidazol-1-yl); $-CH_2-NH-(CH_2)_3$ -(morpholin-4-
 yl); $-CH_2-NH-(CH_2)_4-NHC(NH)NH_2$; $-CH_2-N$ -(2-amino-2-deoxygluconic acid);-

- CH₂-NH-CH₂CH₂-NH-(CH₂)₁₁CH₃; -CH₂-NH-CH(COOH)CH₂COOH; -CH₂-NH-CH₂CH₂-NHSO₂-(CH₂)₇CH₃; -CH₂-NH-CH₂CH₂-NHSO₂-(CH₂)₈CH₃; -CH₂-NH-CH₂CH₂-NHSO₂-(CH₂)₉CH₃; -CH₂-NH-CH₂CH₂-NHSO₂-(CH₂)₁₁CH₃; -CH₂-NH-CH₂CH₂-NH-(CH₂)₇CH₃; -CH₂-NH-CH₂CH₂-O-CH₂CH₂OH; -CH₂-NH-CH₂CH₂C(O)-N-(D-glucosamine); -CH₂-NH-(6-oxo-[1,3]oxazinan-3-yl); -CH₂-NH-CH₂CH₂-S-(CH₂)₇CH₃; -CH₂-NH-CH₂CH₂-S-(CH₂)₈CH₃; -CH₂-NH-CH₂CH₂-S-(CH₂)₉CH₃; -CH₂-NH-CH₂CH₂-S-(CH₂)₁₁CH₃; -CH₂-NH-CH₂CH₂-S-(CH₂)₆Ph; -CH₂-NH-CH₂CH₂-S-(CH₂)₈Ph; -CH₂-NH-CH₂CH₂-S-(CH₂)₁₀Ph; -CH₂-NH-CH₂CH₂-S-CH₂-(4-(4-CF₃-Ph)Ph); -CH₂-NH-CH₂CH₂-NH-(CH₂)₁₁CH₃;
- 10 or -CH₂-NH-(CH₂)₅-COOH.

Preferably, R⁸ is -CH₂C(O)NH₂, -CH₂COOH, benzyl, 4-hydroxyphenyl or 3-chloro-4-hydroxyphenyl.

Preferably, R⁹ is hydrogen or alkyl.

- Preferably, R¹⁰ is alkyl or substituted alkyl. More preferably, R¹⁰ is the
- 15 side-chain of a naturally-occurring amino acid, such as isobutyl.

Preferably, R¹¹ is hydrogen or alkyl.

- Preferably, R¹² is hydrogen, alkyl, substituted alkyl or -C(O)R^d. R¹² can also preferably be hydrogen; -CH₂COOH; -CH₂-[CH(OH)]₅CH₂OH; -CH₂CH(OH)CH₂OH; -CH₂CH₂NH₂; -CH₂C(O)OCH₂CH₃; -CH₂-(2-pyridyl); -CH₂-[CH(OH)]₄COOH; -CH₂-(3-carboxyphenyl); (R)-C(O)CH(NH₂)(CH₂)₄NH₂; -C(O)Ph; -C(O)CH₂NHC(O)CH₃; E-CH₂CH₂-S-(CH₂)₃CH=CH(CH₂)₄CH₃; or -C(O)CH₃.
- 20


Preferably, X¹ and X² are each chloro.

Preferably, X³ is hydrogen.

- 25 Preferably, each Y is independently selected from the group consisting of oxygen, sulfur, -S-S-, -NR^c-, -S(O)-, -SO₂-, -NR^cC(O)-, -OSO₂-, -OC(O)-, -NR^cSO₂-, -C(O)NR^c-, -C(O)O-, -SO₂NR^c-, -SO₂O-, -P(O)(OR^c)O-, -P(O)(OR^c)NR^c-, -OP(O)(OR^c)O-, -OP(O)(OR^c)NR^c-, -OC(O)O-, -NR^cC(O)O-, -NR^cC(O)NR^c-, -OC(O)NR^c-, and
- 30 -NR^cSO₂NR^c-;

Preferably, n is 0 or 1, and more preferably, n is 1.

Another preferred compound of the invention is a glycopeptide of formula II:



wherein:

R^{19} is hydrogen;

- 5 R^{20} is $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$; and R^a , Y , R^b , Z , x , R^f , R^3 , and R^5 have any of the values or preferred values described herein;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;

provided at least one of R^3 and R^5 is a substituent that comprises a

- 10 saccharide group and a carboxy group.

A preferred value for R^{15} , R^{20} , or $-R^a-Y-R^b-(Z)_x$ is

- CH₂CH₂-NH-(CH₂)₉CH₃; -CH₂CH₂CH₂-NH-(CH₂)₈CH₃;
 -CH₂CH₂CH₂CH₂-NH-(CH₂)₇CH₃; -CH₂CH₂-NH-SO₂-(CH₂)₉CH₃;
 -CH₂CH₂-NH-SO₂-(CH₂)₁₁CH₃; -CH₂CH₂-S-(CH₂)₈CH₃;
 15 -CH₂CH₂-S-(CH₂)₉CH₃; -CH₂CH₂-S-(CH₂)₁₀CH₃;
 -CH₂CH₂CH₂-S-(CH₂)₈CH₃; -CH₂CH₂CH₂-S-(CH₂)₉CH₃;
 -CH₂CH₂CH₂-S-(CH₂)₃-CH=CH-(CH₂)₄CH₃ (*trans*); -CH₂CH₂CH₂CH₂-
 S-(CH₂)₇CH₃; -CH₂CH₂-S(O)-(CH₂)₉CH₃; -CH₂CH₂-S-(CH₂)₆Ph;
 -CH₂CH₂-S-(CH₂)₈Ph; -CH₂CH₂CH₂-S-(CH₂)₈Ph; -CH₂CH₂-NH-CH₂-4-(4-
 20 Cl-Ph)-Ph; -CH₂CH₂-NH-CH₂-4-[4-(CH₃)₂CHCH₂]-Ph; -CH₂CH₂-NH-CH₂-

- 4-(4-CF₃-Ph)-Ph; -CH₂CH₂-S-CH₂-4-(4-Cl-Ph)-Ph; -CH₂CH₂-S(O)-CH₂-4-(4-Cl-Ph)-Ph; -CH₂CH₂CH₂-S-CH₂-4-(4-Cl-Ph)-Ph;
 -CH₂CH₂CH₂-S(O)-CH₂-4-(4-Cl-Ph)-Ph; -CH₂CH₂CH₂-S-CH₂-4-[3,4-di-Cl-PhCH₂O-)-Ph; -CH₂CH₂-NHSO₂-CH₂-4-[4-(4-Ph)-Ph]-Ph;
 5 -CH₂CH₂CH₂-NHSO₂-CH₂-4-(4-Cl-Ph)-Ph; -CH₂CH₂CH₂-NHSO₂-CH₂-4-(Ph-C≡C-)-Ph; -CH₂CH₂CH₂-NHSO₂-4-(4-Cl-Ph)-Ph; or
 -CH₂CH₂CH₂-NHSO₂-4-(naphth-2-yl)-Ph. Another preferred value for R¹⁵ or R²⁰ is 4-(4-chlorophenyl)benzyl or 4-(4-chlorobenzyloxy)benzyl.

The invention also provides a pharmaceutical composition comprising a
 10 pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of the invention. In one preferred embodiment, the pharmaceutically acceptable carrier comprises an aqueous cyclodextrin solution. Preferably, the cyclodextrin is hydroxypropyl-β-cyclodextrin or sulfobutyl ether β-cyclodextrin. More preferably, the cyclodextrin is hydroxypropyl-β-cyclodextrin.

15 The compounds of the invention are highly effective antibacterial agents. Accordingly, the invention also provides a method of treating a mammal having a bacterial disease, comprising administering to the mammal a therapeutically effective amount of a compound of the invention. The invention also provides a method of treating a mammal having a bacterial disease, comprising
 20 administering to the mammal a therapeutically effective amount of a pharmaceutical composition of the invention.

The invention also provides processes and intermediates useful for preparing compounds of the invention, which processes and intermediates are described further herein.

25 The invention also provides a compound of the invention as described herein for use in medical therapy, as well as the use of a compound of the invention in the manufacture of a formulation or medicament for treating a bacterial disease in a mammal.

Preferred compounds of the invention are the compounds of formula II
 30 shown in Table I below wherein R¹⁹ is hydrogen.

Table I: Preferred Compounds of formula II

Compound	R ³	R ⁵	R ²⁰
1	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	CH ₃ (CH ₂) ₁₁ -
2	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	CH ₃ (CH ₂) ₁₂ -
3	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ NHCH ₂ CH ₂ -
4	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ SCH ₂ CH ₂ -
5	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ OCH ₂ CH ₂ -
6	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	4-(4-chlorophenyl)-benzyl
7	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-chlorophenyl)-benzylamino)ethyl
8	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-trifluoromethyl-phenyl)benzylamino)ethyl-
9	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	4-(4'-chlorobiphenyl)butyl
10	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-chlorophenyl)-benzylthio)ethyl
11	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-chlorophenyl)-benzyloxy)ethyl
12	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-methyl-benzyloxy)benzylamino)-ethyl-
13	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	2-(4-(4-chloro-benzyloxy)benzylthio)ethyl
14	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	Ph(CH ₂) ₈ S(CH ₂) ₂ NH-
15	Formula III, R ² =N-(D-glucosamine; R ^h =OH	H	Ph(CH ₂) ₈ NH(CH ₂) ₂ -
16	Formula III, R ² =N-(N-methyl-D-glucamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ NHCH ₂ CH ₂ -

Compound	R ³	R ⁵	R ²⁰
17	Formula III, R ^ε =N-(N-methyl-D-glucamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ SCH ₂ CH ₂ -
18	Formula III, R ^ε =N-(N-methyl-D-glucamine; R ^h =OH	H	CH ₃ (CH ₂) ₉ OCH ₂ CH ₂ -
19	Formula III, R ^ε =N-(N-methyl-D-glucamine; R ^h =OH	H	4-(4-chlorophenyl) benzyl
20	OH	Formula V	4-(4-chlorophenyl)-benzyl
21	OH	Formula V	CH ₃ (CH ₂) ₉ SCH ₂ CH ₂ -
22	OH	Formula V	CH ₃ (CH ₂) ₉ NHCH ₂ CH ₂ -
23	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	CH ₃ (CH ₂) ₉ NHCH ₂ CH ₂ -
24	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	CH ₃ (CH ₂) ₉ OCH ₂ CH ₂ -
25	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	CH ₃ (CH ₂) ₉ SCH ₂ CH ₂ -
26	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	4-(4-chlorophenyl) benzyl
27	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	2-(4-(4-trifluoromethyl-phenyl)benzylamino) ethyl
28	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	2-(4-(4-chlorophenyl)-benzylamino)ethyl

Compound	R ³	R ⁵	R ²⁰
29	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	2-(4-(4-methylbenzyloxy)-benzylamino)ethyl
30	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	2-(4-(4-chloro-benzyloxy)benzylthio)-ethyl
31	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	Ph(CH ₂) ₈ NH(CH ₂) ₂ -
32	OH	Formula IV, R ^m =N-(N-methyl-D-glucamine; R ⁿ =OH	Ph(CH ₂) ₈ S(CH ₂) ₂ -

Another preferred group of compounds of the invention are derivatives of the glycopeptide antibiotic A82846B (also known as chloroorienticin A or LY264826; see for example R. Nagarajan et al., *J. Org. Chem.*, **1988**, *54*, 983-986; and N. Tsuji et al., *J. Antibiot.*, **1988**, *41*, 819-822.) that are substituted at the C-terminus with a substituent comprising one or more saccharide groups and a carboxy group, and/or that are substituted at the R-terminus with a substituent that comprises one or more (e.g. 1, 2, 3, 4, or 5) saccharide groups and a carboxy group. The structure of A82846B is similar to vancomycin, except A82846B contains an additional amino sugar (i.e. 4-epi-vancosamine attached at the R² position in formula I.) and further contains 4-epi-vancosamine in place of vancosamine in the disaccharide moiety attached at the R¹ position in formula I.

A preferred group of compounds of the invention are N-alkylated derivatives of A82846B that are substituted at the C-terminus with a substituent that comprises one or more (e.g. 1, 2, 3, 4, or 5) saccharide groups and a carboxy group; or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof. Another preferred group of compounds of the invention are N-alkylated derivatives of A82846B that are substituted at the R-terminus with a substituent that comprises one or more (e.g. 1, 2, 3, 4, or 5) saccharide groups and a carboxy group; or a

pharmaceutically acceptable salt, stereoisomer, or prodrug thereof. A more preferred group of compounds of the invention are C-terminus and/or R-terminus saccharide derivatives of A82846B having a 4-(4-chlorophenyl)benzyl group or a 4-(4-chlorobenzoyloxy)benzyl group attached at the amino group of the 4-epi-
5 vancosamine of the disaccharide moiety. The compounds of the invention that are C-terminus and/or R-terminus saccharide derivatives of A82846B can readily be prepared using the procedures described herein.

The saccharide compounds of the invention have been found to unexpectedly exhibit reduced tissue accumulation and/or nephrotoxicity when
10 administered to a mammal. While not wishing to be bound by theory, it is believed that the saccharide moiety and the carboxy group serve to increase the overall negative charge and the polarity of the glycopeptide under physiological conditions thereby facilitating excretion from the mammal after administration. The unexpected increase in excretion of the saccharide compounds of the
15 invention may be responsible for the reduced tissue accumulation and/or reduced nephrotoxicity observed for these compounds relative to the corresponding compounds that lack the saccharide/carboxy group functionality.

DETAILED DESCRIPTION OF THE INVENTION

20 This invention relates to novel compounds, which are C-terminus or R-terminus saccharide derivatives of glycopeptide antibiotics, as well as to compositions comprising such compounds and to therapeutic methods comprising the administration of such compounds. When describing the compounds, compositions and methods of the invention, the following terms
25 have the following meanings, unless otherwise indicated.

Definitions

The term "alkyl" refers to a monoradical branched or unbranched saturated hydrocarbon chain preferably having from 1 to 40 carbon atoms, more
30 preferably 1 to 10 carbon atoms, and even more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *n*-hexyl, *n*-decyl, tetradecyl, and the like.

The term "substituted alkyl" refers to an alkyl group as defined above, having from 1 to 8 substituents, preferably 1 to 5 substituents, and more preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, guanido, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₃H, and -SO₂-heteroaryl.

The term "alkylene" refers to a diradical of a branched or unbranched saturated hydrocarbon chain, preferably having from 1 to 40 carbon atoms, preferably 1-10 carbon atoms, more preferably 1-6 carbon atoms. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂CH₂-), the propylene isomers (e.g., -CH₂CH₂CH₂- and -CH(CH₃)CH₂-) and the like.

The term "substituted alkylene" refers to an alkylene group, as defined above, having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl. Additionally, such substituted alkylene groups include those where 2 substituents on the alkylene group are fused to form one or more cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heterocyclic or heteroaryl groups fused to the alkylene group. Preferably such fused groups contain from 1 to 3 fused ring structures. Additionally, the term substituted alkylene includes alkylene groups in which from 1 to 5 of the alkylene carbon atoms are replaced with oxygen, sulfur or -NR- where R is hydrogen or alkyl.

Examples of substituted alkenes are chloromethylene (-CH(Cl)-), aminoethylene (-CH(NH₂)CH₂-), 2-carboxypropylene isomers (-CH₂CH(CO₂H)CH₂-), ethoxyethyl (-CH₂CH₂-O-CH₂CH₂-) and the like.

The term "alkaryl" refers to the groups -alkylene-aryl and -substituted alkylene-aryl where alkylene, substituted alkylene and aryl are defined herein. Such alkaryl groups are exemplified by benzyl, phenethyl and the like.

The term "alkoxy" refers to the groups alkyl-O-, alkenyl-O-, cycloalkyl-O-, cycloalkenyl-O-, and alkynyl-O-, where alkyl, alkenyl, cycloalkyl, cycloalkenyl, and alkynyl are as defined herein. Preferred alkoxy groups are alkyl-O- and include, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

The term "substituted alkoxy" refers to the groups substituted alkyl-O-, substituted alkenyl-O-, substituted cycloalkyl-O-, substituted cycloalkenyl-O-, and substituted alkynyl-O- where substituted alkyl, substituted alkenyl, substituted cycloalkyl, substituted cycloalkenyl and substituted alkynyl are as defined herein.

The term "alkylalkoxy" refers to the groups -alkylene-O-alkyl, alkylene-O-substituted alkyl, substituted alkylene-O-alkyl and substituted alkylene-O-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein. Preferred alkylalkoxy groups are alkylene-O-alkyl and include, by way of example, methylenemethoxy (-CH₂OCH₃), ethylenemethoxy (-CH₂CH₂OCH₃), *n*-propylene-*iso*-propoxy (-CH₂CH₂CH₂OCH(CH₃)₂), methylene-*t*-butoxy (-CH₂-O-C(CH₃)₃) and the like.

The term "alkylthioalkoxy" refers to the group -alkylene-S-alkyl, alkylene-S-substituted alkyl, substituted alkylene-S-alkyl and substituted alkylene-S-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein. Preferred alkylthioalkoxy groups are alkylene-S-alkyl and include, by way of example, methylenethiomethoxy (-CH₂SCH₃), ethylenethiomethoxy (-CH₂CH₂SCH₃), *n*-propylene-*iso*-thiopropoxy (-CH₂CH₂CH₂SCH(CH₃)₂), methylene-*t*-thiobutoxy (-CH₂SC(CH₃)₃) and the like.

The term "alkenyl" refers to a monoradical of a branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of vinyl unsaturation.

- 5 Preferred alkenyl groups include ethenyl ($-\text{CH}=\text{CH}_2$), *n*-propenyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), *iso*-propenyl ($-\text{C}(\text{CH}_3)=\text{CH}_2$), and the like.

- The term "substituted alkenyl" refers to an alkenyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.
- 10
15

- The term "alkenylene" refers to a diradical of a branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of vinyl unsaturation. This term is exemplified by groups such as ethenylene ($-\text{CH}=\text{CH}-$), the propenylene isomers (e.g., $-\text{CH}_2\text{CH}=\text{CH}-$ and $-\text{C}(\text{CH}_3)=\text{CH}-$) and the like.
- 20

- The term "substituted alkenylene" refers to an alkenylene group as defined above having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and
- 25
30

-SO₂-heteroaryl. Additionally, such substituted alkenylene groups include those where 2 substituents on the alkenylene group are fused to form one or more cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heterocyclic or heteroaryl groups fused to the alkenylene group.

5 The term "alkynyl" refers to a monoradical of an unsaturated hydrocarbon preferably having from 2 to 40 carbon atoms, more preferably 2 to 20 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of acetylene (triple bond) unsaturation. Preferred alkynyl groups include ethynyl ($-C\equiv CH$), propargyl ($-CH_2C\equiv CH$) and
10 the like.

 The term "substituted alkynyl" refers to an alkynyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino,
15 acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -
20 SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

 The term "alkynylene" refers to a diradical of an unsaturated hydrocarbon preferably having from 2 to 40 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of acetylene (triple bond) unsaturation. Preferred
25 alkynylene groups include ethynylene ($-C\equiv C-$), propargylene ($-CH_2C\equiv C-$) and the like.

 The term "substituted alkynylene" refers to an alkynylene group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino,
30 acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted

thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl

- 5 The term "acyl" refers to the groups HC(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, cycloalkenyl-C(O)-, substituted cycloalkenyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic are as
10 defined herein.

- The term "acylamino" or "aminocarbonyl" refers to the group -C(O)NRR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic or where both R groups are joined to form a heterocyclic group (e.g., morpholino) wherein alkyl, substituted alkyl, aryl, heteroaryl and
15 heterocyclic are as defined herein.

- The term "aminoacyl" refers to the group -NRC(O)R where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined
herein.

- 20 The term "aminoacyloxy" or "alkoxycarbonylamino" refers to the group -NRC(O)OR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

- The term "acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, aryl-C(O)O-, heteroaryl-C(O)O-, and heterocyclic-C(O)O- wherein alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, and heterocyclic are as
25 defined herein.

- The term "aryl" refers to an unsaturated aromatic carbocyclic group of
30 from 6 to 20 carbon atoms having a single ring (e.g., phenyl) or multiple condensed (fused) rings, wherein at least one ring is aromatic (e.g., naphthyl, dihydrophenanthrenyl, fluorenyl, or anthryl). Preferred aryls include phenyl, naphthyl and the like.

Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxy, carboxyalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, aminoacyloxy, oxyacylamino, sulfonamide, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl and trihalomethyl. Preferred aryl substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

The term "aryloxy" refers to the group aryl-O- wherein the aryl group is as defined above including optionally substituted aryl groups as also defined above.

The term "arylene" refers to the diradical derived from aryl (including substituted aryl) as defined above and is exemplified by 1,2-phenylene, 1,3-phenylene, 1,4-phenylene, 1,2-naphthylene and the like.

The term "amino" refers to the group -NH₂.

The term "substituted amino" refers to the group -NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, cycloalkenyl, substituted cycloalkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl and heterocyclic provided that both R's are not hydrogen.

"Amino acid" refers to any of the naturally occurring amino acids (e.g. Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D, L, or DL form. The side chains of naturally occurring amino acids are well known in the art and include, for example, hydrogen (e.g., as in glycine), alkyl (e.g., as in alanine, valine, leucine, isoleucine, proline), substituted alkyl (e.g., as in threonine, serine, methionine, cysteine, aspartic acid, asparagine, glutamic acid, glutamine, arginine, and

lysine), alkaryl (e.g., as in phenylalanine and tryptophan), substituted arylalkyl (e.g., as in tyrosine), and heteroarylalkyl (e.g., as in histidine).

The term "carboxy" refers to -COOH.

The term "C-terminus" as it relates to a glycopeptide is well understood in the art. For example, for a glycopeptide of formula I, the C-terminus is the position substituted by the group R³.

The term "dicarboxy-substituted alkyl" refers to an alkyl group substituted with two carboxy groups. This term includes, by way of example, -CH₂(COOH)CH₂COOH and -CH₂(COOH)CH₂CH₂COOH.

The term "carboxyalkyl" or "alkoxycarbonyl" refers to the groups "-C(O)O-alkyl", "-C(O)O-substituted alkyl", "-C(O)O-cycloalkyl", "-C(O)O-substituted cycloalkyl", "-C(O)O-alkenyl", "-C(O)O-substituted alkenyl", "-C(O)O-alkynyl" and "-C(O)O-substituted alkynyl" where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl and substituted alkynyl are as defined herein.

The term "cycloalkyl" refers to cyclic alkyl groups of from 3 to 20 carbon atoms having a single cyclic ring or multiple condensed rings. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

The term "substituted cycloalkyl" refers to cycloalkyl groups having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

The term "cycloalkenyl" refers to cyclic alkenyl groups of from 4 to 20 carbon atoms having a single cyclic ring and at least one point of internal

unsaturation. Examples of suitable cycloalkenyl groups include, for instance, cyclobut-2-enyl, cyclopent-3-enyl, cyclooct-3-enyl and the like.

The term "substituted cycloalkenyl" refers to cycloalkenyl groups having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

The term "halo" or "halogen" refers to fluoro, chloro, bromo and iodo.

"Haloalkyl" refers to alkyl as defined herein substituted by 1-4 halo groups as defined herein, which may be the same or different. Representative haloalkyl groups include, by way of example, trifluoromethyl, 3-fluorododecyl, 12,12,12-trifluorododecyl, 2-bromooctyl, 3-bromo-6-chloroheptyl, and the like.

The term "heteroaryl" refers to an aromatic group of from 1 to 15 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within at least one ring (if there is more than one ring).

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxy, carboxyalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl and trihalomethyl. Preferred aryl substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy. Such heteroaryl groups can

have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indoliziny or benzothienyl). Preferred heteroaryls include pyridyl, pyrrolyl and furyl.

“Heteroarylalkyl” refers to (heteroaryl)alkyl- where heteroaryl and alkyl are as defined herein. Representative examples include 2-pyridylmethyl and the like.

The term “heteroaryloxy” refers to the group heteroaryl-O-.

The term “heteroarylene” refers to the diradical group derived from heteroaryl (including substituted heteroaryl), as defined above, and is exemplified by the groups 2,6-pyridylene, 2,4-pyridylene, 1,2-quinolinyne, 1,8-quinolinyne, 1,4-benzofuranylene, 2,5-pyridylene, 2,5-indolenyl and the like.

The term “heterocycle” or “heterocyclic” refers to a monoradical saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 40 carbon atoms and from 1 to 10 hetero atoms, preferably 1 to 4 heteroatoms, selected from nitrogen, sulfur, phosphorus, and/or oxygen within the ring.

Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, oxo (O=), and -SO₂-heteroaryl. Such heterocyclic groups can have a single ring or multiple condensed rings. Preferred heterocyclics include morpholino, piperidiny, and the like.

Examples of nitrogen heterocycles and heteroaryls include, but are not limited to, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine,

- isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, morpholino,
- 5 piperidinyl, tetrahydrofuranyl, and the like as well as N-alkoxy-nitrogen containing heterocycles.

Another class of heterocyclics is known as "crown compounds" which refers to a specific class of heterocyclic compounds having one or more repeating units of the formula $[-(\text{CH}_2)_a\text{A}-]$ where a is equal to or greater than 2, and A at each separate occurrence can be O, N, S or P. Examples of crown

10 compounds include, by way of example only, $[-(\text{CH}_2)_3\text{-NH-}]_3$, $[-((\text{CH}_2)_2\text{-O})_4\text{-}((\text{CH}_2)_2\text{-NH})_2]$ and the like. Typically such crown compounds can have from 4 to 10 heteroatoms and 8 to 40 carbon atoms.

The term "heterocyclooxy" refers to the group heterocyclic-O-.

- 15 The term "thioheterocyclooxy" refers to the group heterocyclic-S-.

The term "oxyacylamino" or "aminocarbonyloxy" refers to the group $-\text{OC}(\text{O})\text{NRR}$ where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

- 20 The term "prodrug" is well understood in the art and includes compounds that are converted to pharmaceutically active compounds of the invention in a mammalian system. For example, see *Remington's Pharmaceutical Sciences*, 1980, vol. 16, Mack Publishing Company, Easton, Pennsylvania, 61 and 424.

The term "saccharide group" refers to an oxidized, reduced or substituted

25 saccharide monoradical covalently attached to the glycopeptide or other compound via any atom of the saccharide moiety, for example, via the aglycone carbon atom. The term includes amino-containing saccharide groups. Representative saccharides include, by way of illustration, hexoses such as D-glucose, D-mannose, D-xylose, D-galactose, vancosamine, 3-desmethyl-

30 vancosamine, 3-epi-vancosamine, 4-epi-vancosamine, acosamine, actinosamine, daunosamine, 3-epi-daunosamine, ristosamine, D-glucamine, N-methyl-D-glucamine, D-glucuronic acid, N-acetyl-D-glucosamine, N-acetyl-D-galactosamine, sialic acid, iduronic acid, L-fucose, and the like; pentoses such

as D-ribose or D-arabinose; ketoses such as D-ribulose or D-fructose;
disaccharides such as 2-O-(α -L-vancosaminyl)- β -D-glucopyranose, 2-O-(3-
desmethyl- α -L-vancosaminyl)- β -D-glucopyranose, sucrose, lactose, or maltose;
derivatives such as acetals, amines, acylated, sulfated and phosphorylated sugars;
5 oligosaccharides having from 2 to 10 saccharide units. For the purposes of this
definition, these saccharides are referenced using conventional three letter
nomenclature and the saccharides can be either in their open or preferably in
their pyranose form.

The term "amino-containing saccharide group" or "amino saccharide"
10 refers to a saccharide group having an amino substituent. Representative amino-
containing saccharides include L-vancosamine, 3-desmethyl-vancosamine, 3-epi-
vancosamine, 4-epi-vancosamine, acosamine, actinosamine, daunosamine, 3-epi-
daunosamine, ristosamine, N-methyl-D-glucamine and the like.

The term "spiro-attached cycloalkyl group" refers to a cycloalkyl group
15 attached to another ring via one carbon atom common to both rings.

The term "stereoisomer" as it relates to a given compound is well
understood in the art, and refers another compound having the same molecular
formula, wherein the atoms making up the other compound differ in the way
they are oriented in space, but wherein the atoms in the other compound are like
20 the atoms in the given compound with respect to which atoms are joined to
which other atoms (e.g. an enantiomer, a diastereomer, or a geometric isomer).
See for example, *Morrison and Boyde Organic Chemistry*, 1983, 4th ed., Allyn
and Bacon, Inc., Boston, Mass., page 123

The term "sulfonamide" refers to a group of the formula $-\text{SO}_2\text{NRR}$,
25 where each R is independently hydrogen, alkyl, substituted alkyl, aryl,
heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and
heterocyclic are as defined herein.

The term "thiol" refers to the group -SH.

The term "thioalkoxy" refers to the group -S-alkyl.

30 The term "substituted thioalkoxy" refers to the group -S-substituted alkyl.

The term "thioaryloxy" refers to the group aryl-S- wherein the aryl group
is as defined above including optionally substituted aryl groups also defined
above.

The term "thioheteroaryloxy" refers to the group heteroaryl-S- wherein the heteroaryl group is as defined above including optionally substituted aryl groups as also defined above.

The term "thioether derivatives" when used to refer to the glycopeptide
5 compounds of this invention includes thioethers (-S-), sulfoxides (-SO-) and sulfones (-SO₂-).


As to any of the above groups which contain one or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-
10 feasible. In addition, the compounds of this invention include all stereochemical isomers arising from the substitution of these compounds.

"Cyclodextrin" refers to cyclic molecules containing six or more α -D-glucopyranose units linked at the 1,4 positions by α linkages as in amylose. β -Cyclodextrin or cycloheptaamylose contains seven α -D-glucopyranose units. As
15 used herein, the term "cyclodextrin" also includes cyclodextrin derivatives such as hydroxypropyl and sulfobutyl ether cyclodextrins. Such derivatives are described for example, in U.S. Patent Nos. 4,727,064 and 5,376,645. One preferred cyclodextrin is hydroxypropyl β -cyclodextrin having a degree of substitution of from about 4.1-5.1 as measured by FTIR. Such a cyclodextrin is
20 available from Cerestar (Hammond, Indiana, USA) under the name Cavitron™ 82003.

"Glycopeptide" refers to oligopeptide (e.g. heptapeptide) antibiotics, characterized by a multi-ring peptide core optionally substituted with saccharide groups, such as vancomycin. Examples of glycopeptides included in this
25 definition may be found in "Glycopeptides Classification, Occurrence, and Discovery", by Raymond C. Rao and Louise W. Crandall, ("Drugs and the Pharmaceutical Sciences" Volume 63, edited by Ramakrishnan Nagarajan, published by Marcel Dekker, Inc.). Additional examples of glycopeptides are disclosed in U.S. Patent Nos. 4,639,433; 4,643,987; 4,497,802; 4,698,327;
30 5,591,714; 5,840,684; and 5,843,889; in EP 0 802 199; EP 0 801 075; EP 0 667 353; WO 97/28812; WO 97/38702; WO 98/52589; WO 98/52592; and in *J. Amer. Chem. Soc.*, 1996, 118, 13107-13108; *J. Amer. Chem. Soc.*, 1997, 119, 12041-12047; and *J. Amer. Chem. Soc.*, 1994, 116, 4573-4590. Representative

- glycopeptides include those identified as A477, A35512, A40926, A41030, A42867, A47934, A80407, A82846, A83850, A84575, AB-65, Actaplanin, Actinoidin, Ardacin, Avoparcin, Azureomycin, Balhimycin, Chloroorientein, Chloropolysporin, Decaplanin, *N*-demethylvancomycin, Eremomycin,
- 5 Galacardin, Helvecardin, Izupeptin, Kibdelin, LL-AM374, Mannopeptin, MM45289, MM47756, MM47761, MM49721, MM47766, MM55260, MM55266, MM55270, MM56597, MM56598, OA-7653, Orenticin, Parvodicin, Ristocetin, Ristomycin, Synmonicin, Teicoplanin, UK-68597, UK-69542, UK-72051, Vancomycin, and the like. The term "glycopeptide" as used herein is
- 10 also intended to include the general class of peptides disclosed above on which the sugar moiety is absent, i.e. the aglycone series of glycopeptides. For example, removal of the disaccharide moiety appended to the phenol on vancomycin by mild hydrolysis gives vancomycin aglycone. Also within the scope of the invention are glycopeptides that have been further appended with
- 15 additional saccharide residues, especially aminoglycosides, in a manner similar to vancosamine.

"Vancomycin" refers to a glycopeptide antibiotic having the formula:



When describing vancomycin derivatives, the term " N^{van} -" indicates that a substituent is covalently attached to the amino group of the vancosamine moiety

of vancomycin. Similarly, the term "N^{lec}-" indicates that a substituent is covalently attached to the amino group of the leucine moiety of vancomycin.

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes
5 instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted" means that a group may or may not be substituted with the described substituent.

As used herein, the terms "inert organic solvent" or "inert solvent" or "inert diluent" mean a solvent or diluent which is essentially inert under the
10 conditions of the reaction in which it is employed as a solvent or diluent. Representative examples of materials which may be used as inert solvents or diluents include, by way of illustration, benzene, toluene, acetonitrile, tetrahydrofuran ("THF"), dimethylformamide ("DMF"), chloroform ("CHCl₃"), methylene chloride (or dichloromethane or "CH₂Cl₂"), diethyl ether, ethyl
15 acetate, acetone, methylethyl ketone, methanol, ethanol, propanol, isopropanol, tert-butanol, dioxane, pyridine, and the like. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert solvents.

The term "nitrogen-linked" or "N-linked" means a group or substituent is attached to the remainder of a compound (e.g. a compound of formula I) through
20 a bond to a nitrogen of the group or substituent. The term "oxygen-linked" or "O-linked" means a group or substituent is attached to the remainder of a compound (e.g. a compound of formula I) through a bond to an oxygen of the group or substituent. The term "sulfur-linked" means a group or substituent is attached to the remainder of a compound (e.g. a compound of formula I) through
25 a bond to a sulfur of the group or substituent.

"Pharmaceutically acceptable salt" means those salts which retain the biological effectiveness and properties of the parent compounds and which are not biologically or otherwise harmful as the dosage administered. The compounds of this invention are capable of forming both acid and base salts by
30 virtue of the presence of amino and carboxy groups respectively.

Pharmaceutically acceptable base addition salts may be prepared from inorganic and organic bases. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, and

magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines, substituted amines including naturally-occurring substituted amines, and cyclic amines, including isopropylamine, trimethyl amine, diethylamine, triethylamine, tripropylamine, 5 ethanolamine, 2-dimethylaminoethanol, tromethamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, N-alkylglucamines, theobromine, purines, piperazine, piperidine, and N-ethylpiperidine. It should also be understood that other carboxylic acid derivatives would be useful in the practice of this invention, for example 10 carboxylic acid amides, including carboxamides, lower alkyl carboxamides, di(lower alkyl) carboxamides, and the like.

Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids. Salts derived from inorganic acids include hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid 15 and the like. Salts derived from organic acids include acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

20 The compounds of this invention typically contain one or more chiral centers. Accordingly, this invention is intended to include racemic mixtures, diastereomers, enantiomers and mixture enriched in one or more stereoisomer. The scope of the invention as described and claimed encompasses the racemic forms of the compounds as well as the individual enantiomers and non-racemic 25 mixtures thereof.

The term "treatment" as used herein includes any treatment of a condition or disease in an animal, particularly a mammal, more particularly a human, and includes:

- (i) preventing the disease or condition from occurring in a subject 30 which may be predisposed to the disease but has not yet been diagnosed as having it;
- (ii) inhibiting the disease or condition, i.e. arresting its development; relieving the disease or condition, i.e. causing regression of the condition; or.

relieving the conditions caused by the disease, i.e. symptoms of the disease.

The term "disease state which is alleviated by treatment with a broad spectrum antibacterial" or "bacterial disease" as used herein is intended to cover all disease states which are generally acknowledged in the art to be usefully
5 treated with a broad spectrum antibacterial in general, and those disease states which have been found to be usefully treated by the specific antibacterials of this invention. Such disease states include, but are not limited to, treatment of a mammal afflicted with pathogenic bacteria, in particular staphylococci (methicillin sensitive and resistant), streptococci (penicillin sensitive and
10 resistant), enterococci (vancomycin sensitive and resistant), and *Clostridium difficile*.

The term "therapeutically effective amount" refers to that amount which is sufficient to effect treatment, as defined herein, when administered to a mammal in need of such treatment. The therapeutically effective amount will
15 vary depending on the subject and disease state being treated, the severity of the affliction and the manner of administration, and may be determined routinely by one of ordinary skill in the art.

The term "protecting group" or "blocking group" refers to any group which, when bound to one or more hydroxyl, thiol, amino, carboxy or other
20 groups of the compounds, prevents undesired reactions from occurring at these groups and which protecting group can be removed by conventional chemical or enzymatic steps to reestablish the hydroxyl, thio, amino, carboxy or other group. The particular removable blocking group employed is not critical and preferred removable hydroxyl blocking groups include conventional substituents such as
25 allyl, benzyl, acetyl, chloroacetyl, thiobenzyl, benzylidene, phenacyl, t-butyl-diphenylsilyl and any other group that can be introduced chemically onto a hydroxyl functionality and later selectively removed either by chemical or enzymatic methods in mild conditions compatible with the nature of the product. Protecting groups are disclosed in more detail in T.W. Greene and P.G.M. Wuts,
30 "Protective Groups in Organic Synthesis" 3rd Ed., 1999, John Wiley and Sons, N.Y.

Preferred removable amino blocking groups include conventional substituents such as t-butyloxycarbonyl (t-BOC), benzyloxycarbonyl (CBZ),

fluorenylmethoxycarbonyl (FMOC), allyloxycarbonyl (ALOC) and the like, which can be removed by conventional conditions compatible with the nature of the product.


Preferred carboxy protecting groups include esters such as methyl, ethyl, propyl, *t*-butyl etc. which can be removed by mild conditions compatible with the nature of the product.

General Synthetic Procedures

The glycopeptide compounds of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection are well known in the art. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.


In the following reaction schemes, the glycopeptide compounds are depicted in a simplified form as a box "G" that shows the carboxy terminus labeled [C], the vancosamine amino terminus labeled [V], the "non-saccharide" amino terminus (leucine amine moiety) labeled [N], and optionally, the resorcinol moiety labeled [R] as follows:



A glycopeptide compound of the present invention, which is substituted at the C-terminus with a substituent that comprises one or more (e.g. 1, 2, 3, 4, or 5) saccharide groups and a carboxy group, can be prepared by coupling a
 5 corresponding glycopeptide compound wherein the C-terminus is a carboxy group with a suitable protected compound. For example, a glycopeptide compound wherein the C-terminus is a carboxy group can be coupled with a saccharide containing amine, alcohol, or thiol compound to form an amide, an ester, or a thioester, respectively. For example a glycopeptide compound of
 10 formula I wherein R³ is a nitrogen linked substituent that comprises a saccharide and a carboxy group can be prepared by coupling a corresponding glycopeptide compound of formula I wherein R³ is hydroxy with the requisite amino-saccharide/protected-carboxy compound to form a compound of formula I wherein R³ is a nitrogen linked substituent that comprises saccharide and a
 15 protected carboxy group. Subsequent deprotection provides a compound of the invention.

A glycopeptide compound of the present invention, which is substituted at the C-terminus with a substituent that comprises one or more saccharide groups and a carboxy group, and wherein the vancosamine amino terminus (V) is
 20 substituted, can be prepared by first reductively alkylating the corresponding glycopeptide compound wherein the vancosamine amino terminus (V) is the free amine (NH₂) and then coupling the corresponding glycopeptide compound with the requisite compound to provide the compound of formula I.

By way of illustration, a glycopeptide compound, such as vancomycin,
 25 can first be reductive alkylated as shown in the following reaction:



- where A represents R^a minus one carbon atom and R^a, R^b, Y, Z and x are as defined herein. This reaction is typically conducted by first contacting one equivalent of the glycopeptide, i.e., vancomycin, with an excess, preferably
- 5 from 1.1 to 1.3 equivalents, of the desired aldehyde in the presence of an excess, preferably about 2.0 equivalents, of a tertiary amine, such as diisopropylethylamine (DIPEA) and the like. This reaction is typically conducted in an inert diluent, such as DMF or acetonitrile/water, at ambient temperature for about 0.25 to 2 hours until formation of the corresponding imine
- 10 and/or hemiaminal is substantially complete. The resulting imine and/or hemiaminal is typically not isolated, but is reacted *in situ* with a metal hydride reducing agent, such as sodium cyanoborohydride and the like, to afford the corresponding amine. This reaction is preferably conducted by contacting the imine and/or hemiaminal with an excess, preferably about 3 equivalents, of
- 15 trifluoroacetic acid, followed by about 1 to 1.2 equivalents of the reducing agent at ambient temperature in methanol or acetonitrile/water. The resulting alkylated product is readily purified by conventional procedures, such as precipitation and/or reverse-phase HPLC. Surprisingly, by forming the imine and/or hemiaminal in the presence of a trialkyl amine, and then acidifying with
- 20 trifluoroacetic acid before contact with the reducing agent, the selectivity for the reductive alkylation reaction is greatly improved, i.e., reductive alkylation at the amino group of the saccharide (e.g., vancosamine) is favored over reductive alkylation at the N-terminus (e.g., the leucinyl group) by at least 10:1, more preferably 20:1.
- 25 The above process is a significantly improvement over previous methods for selectively alkylating an amino saccharide group of a glycopeptide antibiotic.

Thus, the present invention also provides a method for alkylating a glycopeptide that comprises a saccharide-amine comprising:

combining an aldehyde or ketone, a suitable base, and the glycopeptide, to provide a reaction mixture;

5 acidifying the reaction mixture; and

combining the reaction mixture with a suitable reducing agent, to provide a glycopeptide that is alkylated at the saccharide-amine. Preferably, the glycopeptide comprises at least one amino group other than the saccharide-amine.

10 Preferably, the reductive alkylation at the saccharide-amine is favored over reductive alkylation at another amino group of the glycopeptide by at least about 10:1; and more preferably, by at least about 15:1 or about 20:1.

The reductive alkylation process of the invention is typically carried out in the presence of a suitable solvent or combination of solvents, such as, for
15 example, a halogenated hydrocarbon (e.g. methylene chloride), a linear or branched ether (e.g. diethyl ether, tetrahydrofuran), an aromatic hydrocarbon (e.g. benzene or toluene), an alcohol (methanol, ethanol, or isopropanol), dimethylsulfoxide (DMSO), N,N-dimethylformamide, acetonitrile, water, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, tetramethyl urea, N,N-
20 dimethylacetamide, diethylformamide (DMF), 1-methyl-2-pyrrolidinone, tetramethylenesulfoxide, glycerol, ethyl acetate, isopropyl acetate, N,N-dimethylpropylene urea (DMPU) or dioxane. Preferably the alkylation is carried out in acetonitrile/water, or DMF/methanol.

Preferably the reduction (i.e. treatment with the reducing agent) is carried
25 out in the presence of a protic solvent, such as, for example, an alcohol (e.g. methanol, ethanol, propanol, isopropanol, or butanol), water, or the like.

The reductive alkylation process of the invention can be carried out at any suitable temperature from the freezing point to the reflux temperature of the reaction mixture. Preferably the reaction is carried out at a temperature in the
30 range of about 0 °C to about 100 °C. More preferably at a temperature in a range of about 0 °C to about 50 °C, or in a range of about 20 °C to about 30 °C.

Any suitable base can be employed in the reductive alkylation process of the invention. Suitable bases include tertiary amines (e.g. diisopropylethylamine, N-methylmorpholine or triethylamine) and the like.

Any suitable acid can be used to acidify the reaction mixture. Suitable acids include carboxylic acids (e.g. acetic acid, trichloroacetic acid, citric acid, formic acid, or trifluoroacetic acid), mineral acids (e.g. hydrochloric acid, sulfuric acid, or phosphoric acid), and the like. A preferred acid is trifluoroacetic acid.

Suitable reducing agents for carrying out reductive alkylation process of the invention are known in the art. Any suitable reducing agent can be employed in the methods of the invention, provided it is compatible with the functionality present in the glycopeptide. For example, suitable reducing agents include sodium cyanoborohydride, triacetoxyborohydride, pyridine/borane, sodium borohydride, and zinc borohydride. The reduction can also be carried out in the presence of a transition metal catalyst (e.g. palladium or platinum) in the presence of a hydrogen source (e.g. hydrogen gas or cyclohexadiene). See for example, , *Advanced Organic Chemistry*, Fourth Edition, John Wiley & Sons, New York (1992), 899-900.

The glycopeptide resulting from the reductive alkylation is then coupled with an amine that comprises one or more saccharide groups and a protected-carboxy group. In this reaction, the glycopeptide derivative is typically contacted with the amine in the presence of a peptide coupling reagent, such as PyBOP and HOBt, to provide the amide. This reaction is typically conducted in an inert diluent, such as DMF, at a temperature ranging from about 0°C to about 60°C for about 1 to 24 hours or until the coupling reaction is substantially complete. Subsequent deprotection using conventional procedures and reagents affords the compound of this invention.


If desired, the amine coupling step described above can be conducted first to provide an amide, followed by reductive alkylation and deprotection to afford the compound of the invention.

If desired, the glycopeptide compounds of this invention can also be prepared in a step-wise manner in which a precursor to the $-R^a-Y-R^b-(Z)_x$ group is first attached the glycopeptide by reductive alkylation, followed by subsequent

elaboration of the attached precursor using conventional reagent and procedures to form the $-R^a-Y-R^b-(Z)_x$ group. Additionally, ketones may also be employed in the above-described reductive alkylation reactions to afford α -substituted amines.

- 5 Any glycopeptide having an amino group may be employed in these reductive alkylation reactions. Such glycopeptides are well-known in the art and are either commercially available or may be isolated using conventional procedures. Suitable glycopeptides are disclosed, by way of example, in U.S. Patent Nos. 3,067,099; 3,338,786; 3,803,306; 3,928,571; 3,952,095; 4,029,769; 10 4,051,237; 4,064,233; 4,122,168; 4,239,751; 4,303,646; 4,322,343; 4,378,348; 4,497,802; 4,504,467; 4,542,018; 4,547,488; 4,548,925; 4,548,974; 4,552,701; 4,558,008; 4,639,433; 4,643,987; 4,661,470; 4,694,069; 4,698,327; 4,782,042; 4,914,187; 4,935,238; 4,946,941; 4,994,555; 4,996,148; 5,187,082; 5,192,742; 5,312,738; 5,451,570; 5,591,714; 5,721,208; 5,750,509; 5,840,684; and 15 5,843,889. Preferably, the glycopeptide employed in the above reaction is vancomycin.

- As illustrated in the following scheme, a saccharide/carboxy containing aminoalkyl sidechain at the resorcinol moiety of a glycopeptide, such as vancomycin, can be introduced via a Mannich reaction (in this scheme, the 20 resorcinol moiety is shown for clarity). In this reaction, an amine (NHR^cR^c) and an aldehyde (CH_2O), such as formalin (a source of formaldehyde), are reacted with the glycopeptide under basic conditions to give the glycopeptide derivative:



wherein NR^cR^c together comprises a saccharide group and a carboxy group.

- Compounds of the invention comprising a sulfoxide or sulfone can be 25 prepared from the corresponding thio compounds using conventional reagents and procedures. Suitable reagents for oxidizing a thio compound to a sulfoxide

include, by way of example, hydrogen peroxide, peracides such as 3-chloroperoxybenzoic acid (MCPBA), sodium periodate, sodium chlorite, sodium hypochlorite, calcium hypochlorite, tert-butyl hypochlorite and the like. Chiral oxidizing reagents, (optically active reagents) may also be employed to provide
5 chiral sulfoxides. Such optically active reagents are well-known in the art and include, for example, the reagents described in Kagen et al., *Synlett.*, 1990, 643-650.

The aldehydes and ketones employed in the above reactive alkylation reactions are also well-known in the art and are either commercially available or
10 can be prepared by conventional procedures using commercially available starting materials and conventional reagents (for example see March, *Advanced Organic Chemistry*, Fourth Edition, John Wiley & Sons, New York (1992), and references cited therein).

Polycarboxy amino compounds that are useful for preparing the
15 substituted glycopeptides of the invention are commercially available, or can be prepared using techniques that are known in the art. For example, L-aspartic acid, D-aspartic acid, DL-aspartic acid, N-methyl-D-aspartic acid, L-glutamic acid, D-glutamic acid, D,L-glutamic acid, DL-2-methylglutamic acid, DL-2-aminoadipic acid, D-2-aminoadipic acid, L-2-aminoadipic acid, 3-aminoadipic
20 acid, 2,6-diaminopimelic acid, L-gamma-carboxyglutamic acid, lanthionine, D-cystine, L-cystine, iminodiacetic acid, ethylenediamine-N,N'-diacetic acid, and kainic acid are available from Aldrich Chemical Company, Milwaukee, Wisconsin.

Additional details and other methods for preparing the compounds of this
25 invention are described in the Examples below.

Pharmaceutical Compositions

This invention also includes pharmaceutical composition containing the novel glycopeptide compounds of this invention. Accordingly, the glycopeptide
30 compound, preferably in the form of a pharmaceutically acceptable salt, can be formulated for oral or parenteral administration for the therapeutic or prophylactic treatment of bacterial infections.

By way of illustration, the glycopeptide compound can be admixed with conventional pharmaceutical carriers and excipients and used in the form of tablets, capsules, elixirs, suspensions, syrups, wafers, and the like. Such pharmaceutical compositions will contain from about 0.1 to about 90% by weight of the active compound, and more generally from about 10 to about 30%. The pharmaceutical compositions may contain common carriers and excipients, such as corn starch or gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride, and alginic acid. Disintegrators commonly used in the formulations of this invention include croscarmellose, microcrystalline cellulose, corn starch, sodium starch glycolate and alginic acid.

A liquid composition will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s), for example ethanol, glycerine, sorbitol, non-aqueous solvent such as polyethylene glycol, oils or water, optionally with a suspending agent, a solubilizing agent (such as a cyclodextrin), preservative, surfactant, wetting agent, flavoring or coloring agent. Alternatively, a liquid formulation can be prepared from a reconstitutable powder.

For example a powder containing active compound, suspending agent, sucrose and a sweetener can be reconstituted with water to form a suspension; and a syrup can be prepared from a powder containing active ingredient, sucrose and a sweetener.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid compositions. Examples of such carriers include magnesium stearate, starch, lactose, sucrose, microcrystalline cellulose and binders, for example polyvinylpyrrolidone. The tablet can also be provided with a color film coating, or color included as part of the carrier(s). In addition, active compound can be formulated in a controlled release dosage form as a tablet comprising a hydrophilic or hydrophobic matrix.

A composition in the form of a capsule can be prepared using routine encapsulation procedures, for example by incorporation of active compound and excipients into a hard gelatin capsule. Alternatively, a semi-solid matrix of active compound and high molecular weight polyethylene glycol can be prepared and filled into a hard gelatin capsule; or a solution of active compound in

polyethylene glycol or a suspension in edible oil, for example liquid paraffin or fractionated coconut oil can be prepared and filled into a soft gelatin capsule.

Tablet binders that can be included are acacia, methylcellulose, sodium carboxymethylcellulose, poly-vinylpyrrolidone (Povidone), hydroxypropyl
5 methylcellulose, sucrose, starch and ethylcellulose. Lubricants that can be used include magnesium stearate or other metallic stearates, stearic acid, silicone fluid, talc, waxes, oils and colloidal silica.

Flavoring agents such as peppermint, oil of wintergreen, cherry flavoring or the like can also be used. Additionally, it may be desirable to add a coloring
10 agent to make the dosage form more attractive in appearance or to help identify the product.

The compounds of the invention and their pharmaceutically acceptable salts that are active when given parenterally can be formulated for intramuscular, intrathecal, or intravenous administration.

15 A typical composition for intra-muscular or intrathecal administration will consist of a suspension or solution of active ingredient in an oil, for example arachis oil or sesame oil. A typical composition for intravenous or intrathecal administration will consist of a sterile isotonic aqueous solution containing, for example active ingredient and dextrose or sodium chloride, or a mixture of
20 dextrose and sodium chloride. Other examples are lactated Ringer's injection, lactated Ringer's plus dextrose injection, Normosol-M and dextrose, Isolyte E, acylated Ringer's injection, and the like. Optionally, a co-solvent, for example, polyethylene glycol; a chelating agent, for example, ethylenediamine tetracetic acid; a solubilizing agent, for example, a cyclodextrin; and an anti-oxidant, for
25 example, sodium metabisulphite, may be included in the formulation. Alternatively, the solution can be freeze dried and then reconstituted with a suitable solvent just prior to administration.

In a preferred embodiment, the glycopeptide derivatives of this invention are formulated in an aqueous solution containing a cyclodextrin. In another
30 preferred embodiment the glycopeptide derivatives of this invention are formulated as a lyophilized powder containing a cyclodextrin or as a sterile powder containing a cyclodextrin. Preferably, the cyclodextrin is hydroxypropyl- β -cyclodextrin or sulfobutyl ether β -cyclodextrin; more preferably, the

cyclodextrin is hydroxypropyl- β -cyclodextrin. Typically, in an injectable solution, the cyclodextrin will comprise about 1 to 25 weight percent; preferably, about 2 to 10 weight percent; more preferable, about 4 to 6 weight percent, of the formulation. Additionally, the weight ratio of the cyclodextrin to the
5 glycopeptide derivative will preferably be from about 1:1 to about 10:1.

The compounds of the invention and their pharmaceutically acceptable salts which are active on rectal administration can be formulated as suppositories. A typical suppository formulation will generally consist of active ingredient with a binding and/or lubricating agent such as a gelatin or cocoa
10 butter or other low melting vegetable or synthetic wax or fat.

The compounds of this invention and their pharmaceutically acceptable salts which are active on topical administration can be formulated as transdermal compositions or transdermal delivery devices ("patches"). Such compositions include, for example, a backing, active compound reservoir, a control membrane,
15 liner and contact adhesive. Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991. Such patches may be constructed for
20 continuous, pulsatile, or on demand delivery of pharmaceutical agents.

The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances,
25 including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

Suitable doses are in the general range of from 0.01-100 mg/kg/day, preferably 0.1-50 mg/kg/day. For an average 70 kg human, this would amount
30 to 0.7 mg to 7 g per day, or preferably 7 mg to 3.5 g per day. A more preferred dose for a human is about 500 mg to about 2 g per day.

Other suitable formulations for use in the present invention can be found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

The following formulation examples illustrate representative
5 pharmaceutical compositions of the present invention.

Formulation Example A

This example illustrates the preparation of a representative
pharmaceutical composition for oral administration of a compound of this
10 invention:

	Ingredients	Quantity per tablet, (mg)
	Active Compound	200
	Lactose, spray-dried	148
15	Magnesium stearate	2

The above ingredients are mixed and introduced into a hard-shell gelatin capsule.

Formulation Example B

This example illustrates the preparation of another representative
20 pharmaceutical composition for oral administration of a compound of this
invention:

	Ingredients	Quantity per tablet, (mg)
	Active Compound	400
	Cornstarch	50
	Lactose	145
25	Magnesium stearate	5

30 The above ingredients are mixed intimately and pressed into single scored
tablets.

Formulation Example C

This example illustrates the preparation of a representative
35 pharmaceutical composition for oral administration of a compound of this
invention.

An oral suspension is prepared having the following composition.

Ingredients

	Active Compound	1.0 g
	Fumaric acid	0.5 g
5	Sodium chloride	2.0 g
	Methyl paraben	0.1 g
	Granulated sugar	25.5 g
	Sorbitol (70% solution)	12.85 g
	Veegum K (Vanderbilt Co.)	1.0 g
10	Flavoring	0.035 ml
	Colorings	0.5 mg
	Distilled water	q.s. to 100 mL

15

Formulation Example D

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

An injectable preparation buffered to a pH of 4 is prepared having the following composition:

20

Ingredients

	Active Compound	0.2 g
	Sodium Acetate Buffer Solution (0.4 M)	2.0 mL
25	HCl (1N)	q.s. to pH 4
	Water (distilled, sterile)	q.s. to 20 mL

Formulation Example E

30

This example illustrates the preparation of a representative pharmaceutical composition for injection of a compound of this invention.

A reconstituted solution is prepared by adding 20 mL of sterile water to 1 g of the compound of this invention. Before use, the solution is then diluted with 200 mL of an intravenous fluid that is compatible with the active compound. Such fluids are chosen from 5% dextrose solution, 0.9% sodium chloride, or a mixture of 5% dextrose and 0.9% sodium chloride. Other examples are lactated Ringer's injection, lactated Ringer's plus 5% dextrose injection, Normosol-M and 5% dextrose, Isolyte E, and acylated Ringer's injection.

35

Formulation Example F

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

An injectable preparation is prepared having the following composition:

5

Ingredients

	Active Compound	0.1-5.0 g
	Hydroxypropyl- β -cyclodextrin	1-25 g
10	5% Aqueous Dextrose Solution (sterile)	q.s. to 100 mL

The above ingredients are blended and the pH is adjusted to 3.5 ± 0.5 using 0.5 N HCl or 0.5 N NaOH.

15

Formulation Example G

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

A frozen solution suitable for injection is prepared having the following composition:

20

Frozen Solution

	Active Compound	250 mg to 1000 mg
	Hydroxypropyl- β -cyclodextrin	250 mg to 10 g
25	Excipients - e.g., dextrose	0-50 g
	Water for Injection	10-100 mL

The weight ratio of hydroxypropyl- β -cyclodextrin to the active compound will typically be from about 1:1 to about 10:1.

30

Representative Procedure: Hydroxypropyl- β -cyclodextrin and excipients, if any, are dissolved in about 80% of the water for injection and the active compound is added and dissolved. The pH is adjusted with 1 M sodium hydroxide to 4.7 ± 0.3 and the volume is then adjusted to 95% of the final volume with water for injection. The pH is checked and adjusted, if necessary, and the volume is adjusted to the final volume with water for injection. The formulation is then sterile filtered through a

35

0.22 micron filter and placed into a sterile vial under aseptic conditions.
The vial is capped, labeled and stored frozen.

Formulation Example H

5 This example illustrates the preparation of a representative
pharmaceutical composition containing a compound of this invention.

A lyophilized powder useful for preparing an injectable solution is
prepared having the following composition:

10 Lyophilized Powder

Active Compound	250 mg to 1000 mg
Hydroxypropyl- β -cyclodextrin	250 mg to 10 g
Excipients – e.g., mannitol, sucrose and/or lactose	0-50 g
Buffer agent – e.g., citrate	0-500 mg

The weight ratio of hydroxypropyl- β -cyclodextrin to the active
compound will typically be from about 1:1 to about 10:1.

20 Representative Procedure: Hydroxypropyl- β -cyclodextrin and excipients
and/or buffering agents, if any, are dissolved in about 60% of the water
for injection. The active compound is added and dissolved and the pH is
adjusted with 1 M sodium hydroxide to 4.0-5.0 and the volume is
25 adjusted to 95% of the final volume with water for injection. The pH is
checked and adjusted, if necessary, and the volume is adjusted to the final
volume with water for injection. The formulation is then sterile filtered
through a 0.22 micron filter and placed into a sterile vial under aseptic
conditions. The formulation is then freeze-dried using an appropriate
30 lyophilization cycle. The vial is capped (optionally under partial vacuum
or dry nitrogen), labeled and stored at room temperature or under
refrigeration.

Formulation Example I

35 This example illustrates the preparation of a representative
pharmaceutical composition containing a compound of this invention.

A sterile powder useful for preparing an injectable solution is prepared having the following composition:

5	<u>Sterile Powder</u>	
	Active Compound	250 mg to 1000 mg
	Hydroxypropyl- β -cyclodextrin	250 mg to 10 g ¹
	Excipients	optional
10	The weight ratio of hydroxypropyl- β -cyclodextrin to the active will typically be from about 1:1 to about 10:1.	
15	<u>Representative Procedure:</u> Hydroxypropyl- β -cyclodextrin and the active compound (and any excipients) are dispersed into an appropriate sterile container and the container is sealed (optionally under partial vacuum or dry nitrogen), labeled and stored at room temperature or under refrigeration.	

Administration of representative formulations H and I to a Patient

20 The pharmaceutical formulations described in formulation examples H and I above can be administered intravenously to a patient by the appropriate medical personnel to treat or prevent gram-positive infections. For administration, the above formulations can be reconstituted and/or diluted with a diluent, such as 5% dextrose or sterile saline, as follows:

25 Representative Procedure: The lyophilized powder of formulation example H (e.g., containing 1000 mg of active compound) is reconstituted with 20 mL of sterile water and the resulting solution is further diluted with 80 mL of sterile saline in
30 a 100 mL infusion bag. The diluted solution is then administered to the patient intravenously over 30 to 120 minutes.

Formulation Example I

35 This example illustrates the preparation of a representative pharmaceutical composition for topical application of a compound of this invention.

	Ingredients	grams
	Active compound	0.2-10
	Span 60	2
5	Tween 60	2
	Mineral oil	5
	Petrolatum	10
	Methyl paraben	0.15
	Propyl paraben	0.05
10	BHA (butylated hydroxy anisole)	0.01
	Water	q.s. to 100

All of the above ingredients, except water, are combined and heated to 60°C with stirring. A sufficient quantity of water at 60°C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. 100 g.

Formulation Example K

This example illustrates the preparation of a representative pharmaceutical composition containing a compound of this invention.

A suppository totaling 2.5 grams is prepared having the following composition:

	Ingredients	
25	Active Compound	500 mg
	Witepsol H-15*	balance
	(triglycerides of saturated vegetable fatty acid; a product of Riches-Nelson, Inc., New York, N.Y.)	

Utility

The glycopeptide compounds of this invention, and their pharmaceutically acceptable salts, are useful in medical treatments and exhibit biological activity, including antibacterial activity, which can be demonstrated in using the tests described herein. Such tests are well known to those skilled in the art, and are referenced and described in Lorian "Antibiotics in Laboratory Medicine", Fourth Edition, Williams and Wilkins (1991).

Accordingly, this invention provides methods for treating bacterial or infectious diseases, especially those caused by Gram-positive microorganisms, in

animals. The compounds of this invention are particularly useful in treating infections caused by methicillin-resistant staphylococci. Also, the compounds are useful in treating infection due to enterococci, including vancomycin-resistant enterococci (VRE). Examples of such diseases include severe

5 staphylococcal infections, such as staphylococcal endocarditis and staphylococcal septicemia. The animal treated may be either susceptible to, or infected with, the microorganism. The method of treatment typically comprises administering to the animal an amount of a compound of this invention which is effective for this purpose.

10 In practicing this method, the antibiotic can be administered in a single daily dose or in multiple doses per day. The treatment regimen may require administration over extended periods of time, for example, for several days or for from one to six weeks. The amount per administered dose or the total amount administered will depend on such factors as the nature and severity of

15 the infection, the age and general health of the patient, the tolerance of the patient to the antibiotic and the microorganism or microorganisms in the infection.

Among other properties, the glycopeptide compounds of the invention have been found to have reduced mammalian toxicity when administered to a

20 mammal. For example, the C-terminus and R-terminus saccharide derivatives of the invention have been found to have reduced liver and/or kidney accumulation compared to the corresponding non- saccharide substituted compounds. Moreover, certain compounds of this invention are expected to have reduced nephrotoxicity. Additionally, it has been discovered that the addition of a

25 cyclodextrin compound to a pharmaceutical composition containing the glycopeptide compounds of this invention further reduces the nephrotoxicity and/or tissue accumulation of the glycopeptide compound when administered to a mammal.

The following synthetic and biological examples are offered to illustrate

30 this invention and are not to be construed in any way as limiting the scope of this invention.

EXAMPLES

In the examples below, the following abbreviations have the following meanings. Any abbreviations not defined have their generally accepted meaning. Unless otherwise stated, all temperatures are in degrees Celsius.

5

	ACN	=	acetonitrile
	BOC, Boc	=	<i>tert</i> -butoxycarbonyl
	DIBAL-H	=	diisobutylaluminum hydride
	DIPEA	=	diisopropylethylamine
10	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethyl sulfoxide
	eq.	=	equivalent
	EtOAc	=	ethyl acetate
	Fmoc	=	9-fluorenylmethoxycarbonyl
15	HOBT	=	1-hydroxybenzotriazole hydrate
	Me	=	methyl
	PyBOP	=	benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate
	TEMPO	=	2,2,6,6-tetramethyl-piperidinyloxy, free radical
20	TFA	=	trifluoroacetic acid
	THF	=	tetrahydrofuran
	TLC, tlc	=	thin layer chromatography

In the following examples, vancomycin hydrochloride semi-hydrate was
 25 purchased from Alpharma, Inc. Fort Lee, NJ 07024 (Alpharma AS, Oslo Norway). Other reagents and reactants are available from Aldrich Chemical Co., Milwaukee, WI 53201.


General Procedure A Reductive Alkylation of Vancomycin

30 To a mixture of vancomycin (1 eq.) and the desired aldehyde (1.3 eq.) in DMF was added DIPEA (2 eq.). The reaction was stirred at ambient temperature for 1-2 hours and monitored by reverse-phase HPLC. Methanol and NaCNBH₃ (1 eq.) were added to the solution, followed by TFA (3 eq.). Stirring was continued for an additional hour at ambient temperature. After the reaction was
 35 complete, the methanol was removed *in vacuo*. The residue was precipitated in acetonitrile. Filtration gave the crude product which was then purified by reverse-phase HPLC. If desired, other glycopeptides antibiotics may be used in this procedure.

Example 1**Preparation of Compound 5**

(Formula II wherein R³ is formula III wherein R² is N-(D-glucosamine)

and R^b is OH; R⁵ is hydrogen; R¹⁹ is hydrogen, and R²⁰ is



5


N^{VAN}-Decyloxyethyl vancomycin bistrifluoroacetate (1 g, 0.54 mmol) was dissolved in anhydrous DMF (5 ml) and a solution of PYBOP (0.31 g, 0.59 mmol) and HOBt (0.09 g, 0.59 mmol) in DMF (0.1 mL) was added. After 5 minutes, NMM (0.06 mL, 0.54 mmol) was added, the solution was stirred for 10 20 minutes, and additional NMM (0.06 mL, 0.55 mmol) was added. The solution was stirred for another 5 minutes and a solution of L-glutamic acid δ-N-(D-glucosamine)amide hydrochloride (0.37 g, 1.1 mmol) in DMF (2 mL) and water (0.5 mL) was added, followed immediately by NMM (0.35 mL, 3.2 mmol). The reaction was stirred for 30 minutes, after which it was poured 15 into diethyl ether (140 mL) and acetonitrile (10 mL). The resulting solid was filtered, washed with acetonitrile (2 x 30 mL), dried under reduced pressure, and purified by reverse phase HPLC to give the title compound. MS calculated (M+) 1923.9, found 1925.

20

Example 2**Preparation of Compound 24**

(Formula II wherein R³ is -OH; R⁵ is formula IV wherein R^m is

N-(N-methyl-D-glucamine) and R^a is OH; R¹⁹ is hydrogen, and R²⁰ is



25

N^{VAN}-Decyloxyethyl vancomycin bistrifluoroacetate (0.83 g, 0.45 mmol) and L-glutamic acid δ-N-(N-methyl-D-glucamine)amide hydrochloride (1.6 g, 4.4 mmol) were dissolved in water (7.5 mL) and acetonitrile (7.5 mL). DIPEA (1.17 mL, 6.7 mmol) was added followed by formaldehyde (0.027 mL, 0.36 mmol) diluted in water (0.6 mL) and acetonitrile (0.6 mL). The reaction 30 mixture was stirred at room temperature for 1 hour, The pH of the solution was lowered to 2 with a solution of 50% trifluoroacetic acid in water. The solution was filtered and purified by reverse phase HPLC to give the title compound. MS calculated (M+) 1969.9 found (MH+) 1971.2.

Using the above procedures and the appropriate starting materials the compounds shown in Table I were prepared. The mass spectral data for these compounds were as follows:

5	Compound No.	MW (freebase)	Observed MH ⁺
	1	1907.87	1908.5
	2	1921.89	1923.1
	3	1922.88	1923.8
	4	1939.94	1941.2
10	5	1923.87	1925
	6	1940.21	1940.9
	7	1983.28	1983.9
	8	2016.84	1009.1 (M ₂ +/2)
	9	1982.29	1982.7
15	10	2000.33	2000
	11	1984.27	1984.5
	12	1992.89	1993.5
	13	2030.36	
	14	1987.98	1988.7
20	15	1970.93	1971.5
	16	1938.93	1939.7
	17	1955.98	1956.8
	18	1939.91	1940.9
	19	1956.26	1956.8
25	20	1901.18	1902.1
	21	1900.89	1901.1
	22	1883.85	1884.7
	23	1968.95	1970
	24	1969.94	1971.2
30	25	1986.00	1986.7

5	Compound No.	MW (freebase)	Observed MH ⁺
	26	1986.28	1987.3
	27	2062.91	1032.1 (M2+/2)
	28	2029.35	1015.3 (M2+/2)
	29	2038.96	1020.0 (M2+/2)
5	30	2076.43	
	31	2016.99	1009.1 (M2+/2)
	32	2034.05	1017.8 (M2+/2)

10

Example 3**Determination of Antibacterial Activity****A. *In Vitro* Determination of Antibacterial Activity****1. Determination of Minimal Inhibitory Concentrations (MICs)**

15

Bacterial strains were obtained from either American Type Tissue Culture Collection (ATCC), Stanford University Hospital (SU), Kaiser Permanente Regional Laboratory in Berkeley (KPB), Massachusetts General Hospital (MGH), the Centers for Disease Control (CDC), the San Francisco Veterans' Administration Hospital (SFVA) or the University of California San Francisco Hospital (UCSF). Vancomycin resistant enterococci were phenotyped as Van A or Van B based on their sensitivity to teicoplanin. Some vancomycin resistant enterococci that had been genotyped as Van A, Van B, Van C1 or Van C2 were obtained from the Mayo Clinic.

20

Minimal inhibitory concentrations (MICs) were measured in a microdilution broth procedure under NCCLS guidelines. Routinely, the compounds were serially diluted into Mueller-Hinton broth in 96-well microtiter plates. Overnight cultures of bacterial strains were diluted based on absorbance at 600 nm so that the final concentration in each well was 5×10^5 cfu/mL. Plates were returned to a 35°C incubator. The following day (or 24 hours in the case of Enterococci strains), MICs were determined by visual inspection of the plates. Strains routinely tested in the initial screen included methicillin-sensitive

30

Staphylococcus aureus (MSSA), methicillin-resistant *Staphylococcus aureus*, methicillin-sensitive *Staphylococcus epidermidis* (MSSE), methicillin-resistant *Staphylococcus epidermidis* (MRSE), vancomycin sensitive *Enterococcus faecium* (VSE Fm), vancomycin sensitive *Enterococcus faecalis* (VSE Fs),
5 vancomycin resistant *Enterococcus faecium* also resistant to teicoplanin (VRE Fm Van A), vancomycin resistant *Enterococcus faecium* sensitive to teicoplanin (VRE Fm Van B), vancomycin resistant *Enterococcus faecalis* also resistant to teicoplanin (VRE Fs Van A), vancomycin resistant *Enterococcus faecalis* sensitive to teicoplanin (VRE Fs Van B), enterococcus gallinarum of
10 the Van A genotype (VRE Gm Van A), enterococcus gallinarum of the Van C-1 genotype (VRE Gm Van C-1), enterococcus casseliflavus of the Van C-2 genotype (VRE Cs Van C-2), enterococcus flavescens of the Van C-2 genotype (VRE Fv Van C-2), and penicillin-sensitive *Streptococcus pneumoniae* (PSSP) and penicillin-resistant *Streptococcus pneumoniae* (PSRP). Because of the
15 inability of PSSP and PSRP to grow well in Mueller-Hinton broth, MICs with those strains were determined using either TSA broth supplemented with defibrinated blood or blood agar plates. Compounds which had significant activity against the strains mentioned above were then tested for MIC values in a larger panel of clinical isolates including the species listed above as well as non-
20 speciated coagulase negative *Staphylococcus* both sensitive and resistant to methicillin (MS-CNS and MR-CNS). In addition, they were tested for MICs against gram negative organisms, such as *Escherichia coli* and *Pseudomonas aeruginosa*.

25 2. Determination of Kill Time

Experiments to determine the time required to kill the bacteria were conducted as described in Lorian, "Antibiotics in Laboratory Medicine", Fourth Edition, Williams and Wilkins (1991). These experiments were conducted normally with both staphylococcus and enterococcus strains.

30 Briefly, several colonies were selected from an agar plate and grown at 35°C under constant agitation until it achieved a turbidity of approximately 1.5 and 10⁸ CFU/mL. The sample was then diluted to about 6 x 10⁶ CFU/mL and incubated at 35°C under constant agitation was continued. At various times

aliquots were removed and five ten-fold serial dilutions were performed. The pour plate method was used to determine the number of colony forming units (CFUs).

In general, the compounds of the invention were active in the above tests *in vitro* tests and demonstrated a broad spectrum of activity.

B. In Vivo Determination of Antibacterial Activity

1. Acute Tolerability Studies in Mice

In these studies, a compound of this invention was administered either intravenously or subcutaneously and observed for 5-15 minutes. If there were no adverse effects, the dose was increased in a second group of mice. This dose incrementation continued until mortality occurred, or the dose was maximized. Generally, dosing began at 20 mg/kg and increased by 20 mg/kg each time until the maximum tolerated dose (MTD) is achieved.

2. Bioavailability Studies in Mice

Mice were administered a compound of this invention either intravenously or subcutaneously at a therapeutic dose (in general, approximately 50 mg/kg). Groups of animals were placed in metabolic cages so that urine and feces could be collected for analysis. Groups of animals ($n = 3$) were sacrificed at various times (10 min, 1 hour and 4 hours). Blood was collected by cardiac puncture and the following organs were harvested—lung, liver, heart, brain, kidney, and spleen. Tissues were weighed and prepared for HPLC analysis. HPLC analysis on the tissue homogenates and fluids was used to determine the concentration of the test compound or IiI present. Metabolic products resulting from changes to the test compound were also determined at this juncture.

3. Mouse Septicemia Model

In this model, an appropriately virulent strain of bacteria (most commonly *S. aureus*, or *E. Faecalis* or *E. Faecium*) was administered to mice ($N=5$ to 10 mice per group) intraperitoneally. The bacteria was combined with hog gastric mucin to enhance virulence. The dose of bacteria (normally 10^5 - 10^7) was that sufficient to induce mortality in all of the mice over a three day period. One hour after the

bacteria was administered, a compound of this invention was administered in a single dose either IV or subcutaneously. Each dose was administered to groups of 5 to 10 mice, at doses that typically ranged from a maximum of about 20 mg/kg to a minimum of less than 1 mg/kg. A positive control (normally vancomycin with vancomycin sensitive strains) was administered in each experiment. The dose at which approximately 50% of the animals are saved was calculated from the results.

4. Neutropenic Thigh Model

In this model, antibacterial activity of a compound of this invention was evaluated against an appropriately virulent strain of bacteria (most commonly *S. aureus*, or *E. Faecalis* or *E. Faecium*, sensitive or resistant to vancomycin). Mice were initially rendered neutropenic by administration of cyclophosphamide at 200 mg/kg on day 0 and day 2. On day 4 they were infected in the left anterior thigh by an IM injection of a single dose of bacteria. The mice were then administered the test compound one hour after the bacteria and at various later times (normally 1, 2.5, 4 and 24 hours) the mice were sacrificed (3 per time point) and the thigh excised, homogenized and the number of CFUs (colony forming units) were determined by plating. Blood was also plated to determine the CFUs in the blood.

5. Pharmacokinetic Studies

The rate at which a compound of this invention is removed from the blood can be determined in either rats or mice. In rats, the test animals were cannulated in the jugular vein. The test compound was administered via tail vein injection, and at various time points (normally 5, 15, 30, 60 minutes and 2, 4, 6 and 24 hours) blood was withdrawn from the cannula. In mice, the test compound was also administered via tail vein injection, and at various time points. Blood was normally obtained by cardiac puncture. The concentration of the remaining test compound was determined by HPLC.

In general, the compounds of the invention were active in the above test in vivo and demonstrated a broad spectrum of activity.

Example 4

Determination of Tissue Accumulation

A. Tissue Distribution Using Radiolabeled Compound

This procedure is used to examine the tissue distribution, excretion and metabolism of a radiolabeled test compound in both male and female rats following intravenous infusion at 10 mg/kg. Male and female Sprague-Dawley rats (n = 2 per sex per compound) are dosed with ^3H -labeled test compound at 10 (400 $\mu\text{Ci/kg}$) and 12.5 mg/kg (100 $\mu\text{Ci/kg}$), respectively, via intravenous infusion (~2 min). The test compound is formulated in 5% hydroxypropyl- β -cyclodextrin as 2.5 mg/mL solution. Urine and feces are cage collected over 24 hours period. At 24 hours after dosing, animals are sacrificed and tissues are removed. Serum, urine and tissues are analyzed for total radioactivity by oxidation followed by liquid scintillation counting. Urine and selected tissues samples are extracted and analyzed by reverse phase HPLC with radioactive flow detector for the presence of potential metabolites.

15

B. Tissue Accumulation Following Single Dose

This procedure is used to evaluate tissue distribution of a test compound in rats following single dose administration by infusion. Male Sprague-Dawley rats (n = 3 per dose groups) are dosed with 50 mg/kg of a test compound. Two formulations are used: 30% PEG 400 and 10% sulfobutylether- β -cyclodextrin. Urine samples are cage collected over 24 hours. Blood samples are collected for serum chemistry and concentration determination. Liver and kidneys are removed for histology evaluation. One kidney and part of the liver are homogenized for concentration analysis using reverse phase HPLC with UV detection. Drug concentrations in urine and serum samples are determined by LC-MS analysis.

20

C. Tissue Distribution Following Multiple Doses

This procedure is used too evaluate the potential tissue accumulation of a test compound in rats following multiple dose administration by intravenous infusion. Male and female Sprague-Dawley rats (n = 4 per sex per dose group) are dosed with a test compound at 12.5, 25 and 50 mg/kg per day for seven days. Animals are sacrificed at day 1 (n = 3 per sex per dose group) following the last dose administered. One animal per sex per dose group is retained as recovery animal and


30

sacrificed at day 7 following the last dose administered. The test compound is formulated in 5% hydroxypropyl- β -cyclodextrin or 1% sucrose/4.5% dextrose. Urine samples are cage collected at days 1 and 7 post-dose. Blood samples are collected for serum chemistry and concentration determination. Liver and kidneys
5 are removed for histology evaluation. One kidney and part of the liver are homogenized for concentration analysis using reverse phase HPLC with UV detection. Drug concentrations in urine and serum samples are determined by LC-MS analysis.

While the present invention has been described with reference to the specific
10 embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All
15 such modifications are intended to be within the scope of the claims appended hereto. Additionally, all publications, patents, and patent documents cited hereinabove are incorporated by reference herein in full, as though individually incorporated by reference.


WHAT IS CLAIMED IS:

1. A glycopeptide substituted at the C-terminus and/or the R-terminus with a substituent that comprises one or more saccharide groups and a carboxy group; or a
- 5 pharmaceutically acceptable salt, or stereoisomer, or prodrug thereof;
- provided the glycopeptide is not a compound of formula II:




- a) wherein R³ is N-(2-amino-2-deoxygluconic acid); R⁵ is hydrogen; R¹⁹ is hydrogen; and R²⁰ is -NH-CH₂CH₂-NH-(CH₂)₉CH₃; or
- b) wherein R³ is OH; R⁵ is -CH₂-N-(2-amino-2-deoxygluconic acid); R¹⁹ is hydrogen; and R²⁰ is -CH₂CH₂-NH-(CH₂)₉CH₃.
2. The glycopeptide of claim 1 which is substituted at the C-terminus and the
- 15 R-terminus with a substituent that comprises a saccharide and a carboxy group.
3. The glycopeptide of claim 1 which is substituted at the C-terminus with a
- 20 substituent that comprises a saccharide and a carboxy group.

4. The glycopeptide of claim 1 which is substituted at the R-terminus with a substituent that comprises a saccharide group and a carboxy group.
5. The glycopeptide of claim 3 wherein the substituent that comprises a saccharide and a carboxy group has the formula $-N(R^w)-R^y-R^x$; wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.
6. The glycopeptide of claim 3 wherein the substituent is a substituent of formula III:




wherein one of R^g and R^h is a saccharide, and the other of R^g and R^h is OH.

7. The glycopeptide of claim 4 wherein the substituent that comprises a saccharide and a carboxy group has the formula $-CH_2N(R^w)-R^y-R^x$; wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.
8. The glycopeptide of claim 4 wherein the substituent is a substituent of formula IV:




wherein one of R^m and R^n is a saccharide, and the other is OH.

9. The glycopeptide of claim 4 wherein the substituent is a substituent of formula V:



10. The glycopeptide of claim 1 which is a compound of formula I:



wherein:

R^1 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic and
5 $-R^a-Y-R^b-(Z)_x$; or R^1 is a saccharide group optionally substituted with $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$;

R^2 is hydrogen or a saccharide group optionally substituted with $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$;

10 R^3 is $-OR^c$, $-NR^cR^c$, $-O-R^a-Y-R^b-(Z)_x$, $-NR^c-R^a-Y-R^b-(Z)_x$, $-NR^cR^c$, or $-O-R^c$; or R^3 is a substituent that comprises a saccharide group and a carboxy group;

R^4 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, $-R^a-Y-R^b-(Z)_x$, $-C(O)R^d$

and a saccharide group optionally substituted with $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$.

R^5 is selected from the group consisting of hydrogen, halo, $-\text{CH}(R^c)-\text{NR}^cR^c$, $-\text{CH}(R^c)-\text{NR}^cR^c$, $-\text{CH}(R^c)-\text{NR}^c-R^a-Y-R^b-(Z)_x$, $-\text{CH}(R^c)-R^x$,

- 5 $-\text{CH}(R^c)-\text{NR}^c-R^a-C(=O)-R^x$, or R^5 is substituent that comprises a saccharide group and a carboxy group;

R^6 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, $-R^a-Y-R^b-(Z)_x$, $-C(O)R^d$ and a saccharide group optionally substituted with $-\text{NR}^c-R^a-Y-R^b-(Z)_x$, or R^5 and

- 10 R^6 can be joined, together with the atoms to which they are attached, form a heterocyclic ring optionally substituted with $-\text{NR}^c-R^a-Y-R^b-(Z)_x$;

R^7 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, $-R^a-Y-R^b-(Z)_x$, and $-C(O)R^d$;

- 15 R^8 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

- R^9 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;
- 20

- R^{10} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic; or R^8 and R^{10} are joined to form $-\text{Ar}^1-\text{O}-\text{Ar}^2-$, where Ar^1 and Ar^2 are
- 25 independently arylene or heteroarylene;

- R^{11} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic, or R^{10} and R^{11} are joined, together with the carbon and nitrogen atoms
- 30 to which they are attached, to form a heterocyclic ring;

R^{12} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl,

heterocyclic, $-C(O)R^d$, $-C(NH)R^d$, $-C(O)NR^cR^e$, $-C(O)OR^d$, $-C(NH)NR^cR^e$ and $-R^a-Y-R^b-(Z)_n$, or R^{11} and R^{12} are joined, together with the nitrogen atom to which they are attached, to form a heterocyclic ring;

R^{13} is selected from the group consisting of hydrogen or $-OR^{14}$;

5 R^{14} is selected from hydrogen, $-C(O)R^d$ and a saccharide group;

each R^a is independently selected from the group consisting of alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene;

10 each R^b is independently selected from the group consisting of a covalent bond, alkylene, substituted alkylene, alkenylene, substituted alkenylene, alkynylene and substituted alkynylene, provided R^b is not a covalent bond when Z is hydrogen;

each R^c is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, 15 heteroaryl, heterocyclic and $-C(O)R^d$;

each R^d is independently selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl and heterocyclic;

20 R^e is a saccharide group;

each R^f is independently alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, or heterocyclic;

R^a is N-linked amino saccharide or an N-linked heterocycle;

25 X^1 , X^2 and X^3 are independently selected from hydrogen or chloro;

each Y is independently selected from the group consisting of oxygen, sulfur, $-S-S-$, $-NR^c-$, $-S(O)-$, $-SO_2-$, $-NR^cC(O)-$, $-OSO_2-$, $-OC(O)-$, $-NR^cSO_2-$, $-C(O)NR^c-$, $-C(O)O-$, $-SO_2NR^c-$, $-SO_2O-$, $-P(O)(OR^c)O-$, $-P(O)(OR^c)NR^c-$, $-OP(O)(OR^c)O-$, $-OP(O)(OR^c)NR^c-$, $-OC(O)O-$,

30 $-NR^cC(O)O-$, $-NR^cC(O)NR^c-$, $-OC(O)NR^c-$, $-C(=O)-$, and $-NR^cSO_2NR^c-$;

each Z is independently selected from hydrogen, aryl, cycloalkyl, cycloalkenyl, heteroaryl and heterocyclic;

n is 0, 1 or 2; and

x is 1 or 2;

or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;


provided at least one of R^3 and R^5 is a substituent that comprises a saccharide group and a carboxy group.

5

11. The glycopeptide of claim 10 wherein R^1 is a saccharide group optionally substituted with $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)$.

12. The glycopeptide of claim 10 wherein R^1 is a saccharide group of the

10 formula:




wherein R^{15} is $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$; and R^{16} is hydrogen or methyl.

13. The glycopeptide of claim 10 wherein R^3 has the formula $-N(R^w)-R^y-R^x$;
15 wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.

14. The glycopeptide of claim 10 wherein R^5 is a substituent that comprises a saccharide group and a carboxy group having the formula $-CH_2N(R^w)-R^y-R^x$;
20 wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.

15. The glycopeptide of claim 10 which is a compound of formula II:

wherein:



(II)

R^{19} is hydrogen;


R^{20} is $-R^a-Y-R^b-(Z)_x$, R^f , $-C(O)R^f$, or $-C(O)-R^a-Y-R^b-(Z)_x$; and

R^a , Y , R^b , Z , x , R^f , R^3 , and R^5 have the values defined in claim 10;

- 5 or a pharmaceutically acceptable salt, stereoisomer, or prodrug thereof;
provided at least one of R^3 and R^5 is a substituent that comprises a saccharide group and a carboxy group.

16. The glycopeptide of claim 15 wherein R^3 has the formula $-N(R^w)-R^y-R^x$;
10 wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with a carboxy group; and R^x is a saccharide.


17. The glycopeptide of claim 16 wherein R^3 is of the formula III:



wherein one of R^g and R^h is a saccharide, and the other of R^g and R^h is OH.

18. The glycopeptide of claim 15 wherein R^5 is a substituent that comprises a saccharide group and a carboxy group having the formula $-\text{CH}_2\text{N}(\text{R}^w)-\text{R}^y-\text{R}^x$;
 5 wherein R^w is hydrogen or alkyl; R^y is substituted alkylene, which is substituted with at least one carboxy group; and R^x is a saccharide, and R^5 is -OH.


19. The glycopeptide of claim 18 wherein R^5 is of the formula IV:



wherein one of R^m and R^n is a saccharide, and the other is OH.

10

20. The glycopeptide of claim 15 wherein R^5 is of the formula V:



21. A pharmaceutical composition comprising a pharmaceutically acceptable
 15 carrier and a therapeutically effective amount of a compound of any of claims 1-20.
22. The pharmaceutical composition of Claim 21, wherein the pharmaceutically acceptable carrier comprises an aqueous cyclodextrin solution.

23. A compound as described in any one of claims 1-20 for use in medical therapy.

24. The use of a compound as described in any one of claims 1-20 to prepare a
5 medicament useful for treating a bacterial disease in a mammal.

25. A method of treating a mammal having a bacterial disease, the method comprising administering to the mammal a therapeutically effective amount of a glycopeptide of any one of claims 1, 10, and 15.

10

26. A method of treating a mammal having a bacterial disease, the method comprising administering to the mammal a therapeutically effective amount of a pharmaceutical composition of claim 21.

